DRAINAGE ANALYSIS & EROSION & SEDIMENT CONTROL PLAN

Prepared for:

Paul Thibodeau 76 Young Road Barrington, NH 03825

Land of:

Tax Map 240, Lot 8

Young Road, LLC. 76 Young Road Barrington, NH 03825

Prepared by:

Berry Surveying & Engineering 335 Second Crown Point Road Barrington, NH 03825

Project Number: DB 2022-109

September 19, 2023

Table of Contents

USGS Quadrangle	Location Man:	1:24.000	and 1:10.000
USUS Cudulululu	Location Map.	1.2 I,000 (ania 1.10,000

Watershed Plans: 1:24,000 and two 1:12,000 (CalTopo Based Quad)

Design M	lethod Objectives	Page 2
1.0	Existing Conditions Analysis	Page 3
2.0	Proposed Conditions Analysis	Page 3
3.0	Full Comparative Analysis	Page 4
4.0	Erosion and Sediment Control, BMP's	Page 5
5.0	Conclusion	Page 9

Appendix I - Existing Conditions Analysis

2 Yr- 24 Hr Node Listing 10 Yr -24 Hr. Node Listing 25 Yr-24 Hr. Node Listing 50 Yr-24 Hr. Node Listing 100 Yr-24 Hr. Node Listing 25 Yr-24 Hr. Full Summary

Appendix II - Proposed Conditions Analysis

2 Yr-24 Hr. Node Listing 10 Yr-24 Hr. Node Listing 25 Yr-24 HR. Node Listing 50 Yr-24 Hr. Node Listing 100 Yr-24 Hr. Node Listing 25 Yr-24 Hr. Full Summary

Appendix III - Calculations, Charts, & Graphs

USGS StreamStats Report Extreme Precipitation Tables NCRS USDA Web-soil Report


Enclosed: W-1 Sheet Existing Conditions Watershed Plan Sheets 1-5


W-2 Sheet Post Construction Watershed Plan Sheets 6-10


Erosion & Sediment Control Plan


Map by NH GRANIT Legend State - County ☐ City/Town Sch Lat: 43° 12' 34.213009" N Lon: 71° 2' 4.477153" W Map Scale 1: 10,000 © NH GRANIT, www.granit.unh.edu Map Generated: 9/28/2022 Notes VERMONT CONN. RHODE

Map by NH GRANIT









DESIGN METHOD OBJECTIVES

The owners of Tax Map 240, Lot 8, Young Road, Road, LLC., and Paul Thibodeau, are proposing to develop the property on Young Road. Lot 8 is currently vacant land.

This drainage analysis is somewhat atypical in the total area that is being analyzed and the level of detail that is undertaken in that undertaking. In this case the watershed was looked at in two different formats. The first was by utilizing the USGS StreamStats analysis tool which resulted in the watershed being defined as 1.93 square mile, or 1,235 acres. The second form of analysis utilized the more traditional method outlined below utilizing the USDA SCS TR-20 method in the HydroCAD Stormwater Modeling System and utilizing USDA NRCS Websoil soil definitions. The HydroCAD model delineated a watershed area of 1,263 acres, more conservative by 4%. Although the StreamStats analysis takes into consideration ponding within a watershed and this was not modelled in the HydroCAD environment, neither method, in this case, toke into consideration any impoundment upstream that might result from cross culverts and land development. The watershed area takes into consideration area on both sides of several roadways to include Franklin Pierce Highway, Hale, Sawyer, Robinson, and Mast Drive, and Wildcat Road. There are also no provisions within StreamStats to run a proposed drainage model.

An Existing and Proposed Conditions analysis was conducted for the purpose of estimating the peak rate of stormwater run-off to evaluate the effect of the proposed subdivision. There is one existing drainage discharge point which was identified in the existing analysis and duplicated in the proposed conditions analysis. Evaluating two watershed models we have compared the differences in these rates of peak run-off and surface water volume. Sheet W-1, Existing Conditions Watershed Plan, outlines the characteristics of the site in its existing or pre-construction conditions. The second analysis displays the proposed (post-construction) conditions (See Sheet W-2). The analysis was conducted using data for; 2 Yr-24 Hr (3.05"), 10 Yr-24 Hr (4.60"), 25 Yr-24 Hr (5.81"), 50 Yr-24 Hr (6.94"), and 100 YR-24 Hr (9.91") storm events. Storm event analysis was accomplished using the USDA SCS TR-20 method within the HydroCAD Stormwater Modeling System environment. Rainfall quantities are based on the Extreme Precipitation Table for this location from the Northeast Regional Climate Center / Cornell University (http://precip.eas.cornell.edu).

1.0 Existing Conditions Analysis:

Reference: Sheet W1 - Existing Conditions Watershed Plan (Enclosed)

Existing Conditions Plan

The existing watershed currently consist of mostly vacant land, roadways, and single family home and accessory structures, the soils within the development area are made up of multiple soil types, containing Hydrologic Soil Group (HSG) A, B, C & D including thirty-nine different soil and slope definitions. See the provided USDA NRCS Websoil Soils Report for more information. The land cover types involved are grass land, woods, roofs, and paved or gravel roads.

The land area analyzed consists of 1,263 acres, all evaluated as Subcatchment #1. The point of analysis, Final Reach #1, is just down stream of the Locus Parcel where the wetland and channel flow into the more open water of the Richardson Pond. This pond discharges into the Mallego Brook.

The Time of Concentration for the watershed was calculated as 286 minutes and consisted of twenty-tree segments at a total length of 9,776 linear feet. The watershed area covers land area to the north as far as Church Street and Franklin Pierce Highway, crosses Wildcat Road and Young Road, and touches Beauty Hill Road in the southeast corner, to include a portion of Ross Road. There is one major culvert crossing under Young Road.

The watershed consists of 90.33% pervious area and 9.33% impervious area with the USDS NRCS Weighted Curve Number of 56.

2.0 Proposed Conditions Analysis:

Reference: Sheet W2 - Proposed Conditions Watershed Plan (Enclosed)

Proposed Grading & Drainage Plan

The proposed watershed and analysis are the same as the existing watershed and analysis with the addition of the twenty-three residential lots, including house and driveways.

The watershed consists of 90.06% pervious area and 9.94% impervious area with the USDS NRCS Weighted Curve Number of 56.

3.0 FULL COMPARATIVE ANALYSIS

ANALYSIS COMPONENT: PEAK RATE DISCHARGE (Cubic Feet / Second)

		2 Yr	10 Yr	25 Yr	50 Yr	100 Yr
Final Reach #100	Existing	33.75	141.11	264.31	401.33	583.92
	Proposed	33.75	141.11	264.31	401.33	583.92

ANALYSIS <u>COMPONENT: VOLUME (Acre Feet)</u>

		2 Yr	10 Yr	25 Yr	50 Yr	100 Yr
Final Reach #900	Existing	19.923	76.455	137.379	203.763	292.122
	Proposed	19.923	76.455	137.379	203.763	292.122

USGS StreamStats DISCHARGE (Cubic Feet / Second)

		2 Yr	10 Yr	25 Yr	50 Yr	100 Yr
Final Reach #100	Existing		140	193	237	291

The StreamStats report gives results in percent storm probability. (10-percent AEP flood is equivalent to the 10 YR - 24 HR Storm Event and the 1-percent AEP flood is equivalent to the 100 YR - 24 HR Storm Event.

4.0 EROSION and SEDIMENT CONTROL PLAN & BEST MANAGEMENT PRACTICES (BMP's)

Reference: Proposed Site Plan and Grading Plan

Erosion & Sediment Control Plan Erosion & Sediment Control Details

The proposed site development is protected from erosion and the abutting properties are protected from sediment by the use of Best Management Practices as outlined in the New Hampshire Stormwater Manual, Volume 2, Post-Construction Best Management Practices Selection & Design (December 2008, NHDES & US EPA). Any area disturbed by construction will be re-stabilized within 45 days and abutting properties will not be adversely affected by this development. All swales and drainage structures will be constructed and stabilized prior to having run-off directed to them.

Perimeter Control (Silt Fence / SiltSoxx / Erosion Control Mix Berm)

The plan set demonstrates the location of perimeter sediment control. The Erosion and Sediment Control Details, Sheet E-101, has the specifications for installation and maintenance of the silt fence, Filtrexx mulch filled SiltSoxx (or approved equal), and Erosion Control Mix Berm. There are locations on the site, for example bio-media rain garden protection, where SiltSoxx protection is specified

Vegetated Stabilization

All areas that are disturbed during construction will be stabilized with vegetated material within 30 days of breaking ground. Construction will be managed in such a manner that erosion is prevented and that no abutter's property will be subjected to any siltation, unless otherwise permitted. All areas to be planted with grass for long-term cover will follow the specification and on Sheet E-102 using seeding mixture C, as follows:

Mixture	Pounds per Acre	Pounds per 1,000 Sq. Ft.
Tall Fescue	24	0.55
Creeping Red Fescue	24	0.55
Total	48	1.10

Conservation Mix

Virginia Wild Rye	Native	FACW-
Little Bluestem	Native	FACU
Big Bluestem	Native	FAC
Red Fescue	Native	FACU
Switch Grass	Native	FAC
Partridge Pea	Native	FACU

0.57
FACU-
FAC
FACW
NI
FAC

Conservation Mix to be provided by New England Wetland Plants, Inc., Amherst, MA as outline in their New England Conservation / Wildlife Mix or approved equal. Mix to be applied at a rate of 25 lbs. per acre or one-lb. per 1750 square feet. Ratio of seed is proprietary and substitutions are not allowed.

Conservation Mix will used to stabilize all 2:1 slopes and all land area disturbed within the wetland buffer. (If applicable).

<u>Gravel Wetland Mix:</u> The grass that is planted within a subsurface gravel wetland will be a mix designed for both inundation and dry conditions such as Ernst Seeds, Retention Basin Floor Mix ERNMX-126.

Stabilized Construction Entrance

A temporary gravel construction entrance provides an area where mud can be dislodged from tires before the vehicle leaves the construction site to reduce the amount of mud and sediment transported onto paved municipal and state roads. The stone size for the pad should be 3" coarse aggregate, and the pad itself constructed to a minimum length of 75' for the full width of the access road. The aggregate should be placed at least six inches thick. A plan view and profile are shown on Sheet E-101- Erosion & Sediment Control Detail Plan. (If applicable).

Environmental Dust Control

Dust will be controlled on the site by the use of multiple Best Management Practices. Mulching and temporary seeding will be the first line of protection to be utilized where problems occur. If dust problems are not solved by these applications, the use of water and calcium chloride can be applied. Calcium chloride will be applied at a rate that will keep the surface moist but not cause pollution.

Vegetated Filter Strips / Buffers

Filter strips are areas of land with natural or planted vegetation designed to receive sheet run-off from up gradient development. These natural areas, preferably wooded, are effective in removing sediment and sediment-laden pollutants from such run-off, although their effectiveness is severely diminished when forced to deal with concentrated flow and must therefore be equipped with a level-spreading device. Filter strips should not have a slope exceeding fifteen percent and have a minimum length of seventy-five feet. Each buffer is individually specified and limits are located on the Sediment & Erosion Control Plan

Drainage Swales / Stormwater Conveyance Channels

Drainage swales will be stabilized with vegetation for long term cover as outlined below, and on Sheet E-102 using seed mixture C. As a general rule, velocities in the swale should not exceed 3.0 feet per second for a vegetated swale although velocities as high as 4.5 FPS are allowed under certain soil conditions (If applicable).

Construction Sequence

- 1. Cut and remove trees in construction area only as required.
- 2. Construct and/or install temporary and permanent sediment erosion and detention control facilities as required.
- 3. Erosion, sediment and detention control facility shall be installed & stabilized prior to any earth moving operation & or directing runoff to them.
- 4. Clear, cut and dispose of debris in approved facility.
- 5. Construct temporary culverts as required, or directed.
- 6. Construct roadways for access to desired construction areas. All roads shall be stabilized immediately.
- 7. Install pipe and construction associated appurtenances as required or directed. Install rain gardens. All disturbed areas shall stabilized immediately after grading.
- 8. Begin permanent and temporary seeding and mulching. All cut and fill slopes and disturbed areas shall be seeded or mulched as required, or directed. No area is allowed to be disturbed for a length of time that exceeds 60 days before being stabilized. Daily, or as required. All roadways and parking areas shall be stabilized within 72 hours of

- achieving finished grades. All cut and fill slopes shall be stabilized within 72 hours of achieving finished grades.
- 9. Construct temporary berms, drains ditches, silt fences, sediment traps, etc. mulch and seed as required.
- 10. Inspect and maintain all erosion and sediment control measures during construction.
- 11. Complete permanent seeding and landscaping.
- 12. Remove temporary erosion control measures after seeding areas have established themselves and site improvements are complete.
- 13. Smooth and revegetate all disturbed areas.
- 14. Finish grading and / or paving all roadways.

Temporary Erosion Control Measures

- 1. The smallest practical area of land shall be exposed at any one time.
- 2. Erosion, sediment control measures shall be installed as shown on the plans and at locations as required, or directed by the engineer.
- 3. All disturbed areas shall be returned to original grades and elevations. Disturbed areas shall be loamed with a minimum of 4" of loam and seeded with not less than 1.10 pound of seed per 1,000 square feet (48 pounds per acre) of area.
- 4. Silt fences and other barriers shall be inspected periodically and after every rainstorm during the life of the project. All damaged areas shall be repaired, sediment deposits shall periodically be removed and properly disposed of.
- 5. After all disturbed areas have been stabilized, the temporary erosion control measures are to be removed and the area disturbed by the removal smoothed and re-vegetated.
- 6. Areas must be seeded and mulched within 5 days of final grading, permanently stabilized within 15 days of final grading, or temporarily stabilized within 30 days of initial disturbance of soil.

Inspection and Maintenance Schedule

Perimeter control and Best Management Practices will be inspected during and after storm events to ensure that the practice still has integrity and is not allowing sediment to pass. See the Erosion and Sediment Control Detail Plans for specific inspection and maintenance criteria.

5.0 CONCLUSION

Peak rates of runoff and volume are modeled to be equal in the post-construction analysis / condition, as compared to the pre-construction peak rates of runoff and volume during all storm events.

Respectfully Submitted,

BERRY SURVEYING & ENGINEERING

Christopher R. Berry, SIT 567

Principal, President

Kenneth A. Berry PE, LLS, CPSWQ, CPESC, CESSWI

Principal, VP - Technical Operations

Appendix I - Existing Conditions Analysis


25 Yr - 24 Hr. Full Summary 2 Yr - 24 Hr. Node Listing

10 Yr -24 Hr. Node Listing

25 Yr -24 Hr. Node Listing

50 Yr -24 Hr. Node Listing

100 Yr - 24 Hr. Node Listing



Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 2

Project Notes

CarlsonSurface||
RationalHydrographMethod|2|
ModifiedRational|5|1.00|1.00|1.00|
UnitHydrographMethod|1|

Printed 9/19/2023 Page 3

Area Listing (all nodes)

Area	CN	Description
(acres)		(subcatchment-numbers)
35.478	39	>75% Grass cover, Good HSG A (1S)
21.161	61	>75% Grass cover, Good HSG B (1S)
44.540	74	>75% Grass cover, Good HSG C (1S)
6.322	80	>75% Grass cover, Good HSG D (1S)
5.027	96	Gravel surface, HSG A (1S)
3.384	96	Gravel surface, HSG B (1S)
1.789	96	Gravel surface, HSG C (1S)
0.867	96	Gravel surface, HSG D (1S)
8.794	98	Paved parking HSG A (1S)
2.334	98	Paved parking HSG B (1S)
10.326	98	Paved parking HSG C (1S)
2.495	98	Paved parking HSG D (1S)
3.530	98	Roofs HSG A (1S)
0.981	98	Roofs HSG B (1S)
3.333	98	Roofs HSG C (1S)
0.544	98	Roofs HSG D (1S)
4.356	98	Water Surface HSG A (1S)
12.498	98	Water Surface HSG B (1S)
1.070	98	Water Surface HSG C (1S)
71.867	98	Water Surface HSG D (1S)
442.553	30	Woods, Good HSG A (1S)
259.014	55	Woods, Good HSG B (1S)
170.097	70	Woods, Good HSG C (1S)
150.682	77	Woods, Good HSG D (1S)
1,263.040	56	TOTAL AREA

22-109 W-1 Existing AnalysisPrepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 4

Soil Listing (all nodes)

Area	Soil	Subcatchment
(acres)	Group	Numbers
499.737	HSG A	1S
299.372	HSG B	1S
231.154	HSG C	1S
232.776	HSG D	1S
0.000	Other	
1,263.040		TOTAL AREA

22-109 W-1 Existing AnalysisPrepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 5

Ground Covers (all nodes)

HSG-A (acres)	HSG-B (acres)	HSG-C (acres)	HSG-D (acres)	Other (acres)	Total (acres)	Ground Cover	Subcatchment Numbers
35.478	21.161	44.540	6.322	0.000	107.500	>75% Grass cove	er, Good 1S
5.027	3.384	1.789	0.867	0.000	11.066	Gravel surface	1S
8.794	2.334	10.326	2.495	0.000	23.949	Paved parking	1S
3.530	0.981	3.333	0.544	0.000	8.387	Roofs	1S
4.356	12.498	1.070	71.867	0.000	89.791	Water Surface	1S
442.553	259.014	170.097	150.682	0.000	1,022.346	Woods, Good	1S
499.737	299.372	231.154	232.776	0.000	1,263.040	TOTAL AREA	

Type III 24-hr 25Yr-24Hr Rainfall=5.81"

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023

Page 6

Time span=0.00-24.00 hrs, dt=0.05 hrs, 481 points
Runoff by SCS TR-20 method, UH=SCS, Weighted-CN
Reach routing by Dyn-Stor-Ind method - Pond routing by Dyn-Stor-Ind method

Subcatchment S: 1S Runoff Area=55,018,008

Runoff Area=55,018,008 sf 9.67% Impervious Runoff Depth>1.31" Flow Length=9,776' Tc=286.2 min CN=56 Runoff=264.31 cfs 137.379 af

Reach100R: Final Reach#100

Inflow=264.31 cfs 137.379 af Outflow=264.31 cfs 137.379 af

Total Runoff Area = 1,263.040 ac Runoff Volume = 137.379 af Average Runoff Depth = 1.31" 90.33% Pervious = 1,140.913 ac 9.67% Impervious = 122.127 ac

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023

Page 7

Summary for Subcatchment 1S: 1S

CarlsonPlanXYPos|1151812.3297|261093.3900|

Runoff = 264.31 cfs @ 16.24 hrs, Volume= 137.379 af, Depth> 1.31"

Runoff by SCS TR-20 method, UH=SCS, Weighted-CN, Time Span= 0.00-24.00 hrs, dt= 0.05 hrs Type III 24-hr 25Yr-24Hr Rainfall=5.81"

Area (sf)	CN	Description
153,751	98	Roofs HSG A
42,725	98	Roofs HSG B
145,176	98	Roofs HSG C
23,694	98	Roofs HSG D
1,545,405	39	>75% Grass cover, Good HSG A
921,767	61	>75% Grass cover, Good HSG B
1,940,170	74	>75% Grass cover, Good HSG C
275,367	80	>75% Grass cover, Good HSG D
383,064	98	Paved parking HSG A
101,685	98	Paved parking HSG B
449,781	98	Paved parking HSG C
108,672	98	Paved parking HSG D
19,277,593	30	Woods, Good HSG A
11,282,651	55	Woods, Good HSG B
7,409,442	70	Woods, Good HSG C
6,563,714	77	Woods, Good HSG D
189,750	98	Water Surface HSG A
544,416	98	Water Surface HSG B
46,599	98	Water Surface HSG C
3,130,536	98	Water Surface HSG D
218,969	96	Gravel surface, HSG A
147,414	96	Gravel surface, HSG B
77,919	96	Gravel surface, HSG C
37,748	96	Gravel surface, HSG D
55,018,008	56	Weighted Average
49,698,159		90.33% Pervious Area
5,319,849		9.67% Impervious Area

22-109 W-1 Existing AnalysisPrepared by Berry Surveying & Engineering

HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC


Printed 9/19/2023 Page 8

Tc (min)	Length (feet)	Slope (ft/ft)	Velocity (ft/sec)	Capacity (cfs)	Description
14.2	100	0.0600	0.12		Sheet Flow, Segment #1
					Woods: Light underbrush n= 0.400 P2= 3.05"
0.8	59	0.0639	1.26		Shallow Concentrated Flow, Segment #2
					Woodland Kv= 5.0 fps
0.5	59	0.0679	1.82		Shallow Concentrated Flow, Segment #3
					Short Grass Pasture Kv= 7.0 fps
1.7	128	0.0644	1.27		Shallow Concentrated Flow, Segment #4
					Woodland Kv= 5.0 fps
0.0	13	0.0583	4.90		Shallow Concentrated Flow, Segment #5
					Paved Kv= 20.3 fps
1.5	118	0.0657	1.28		Shallow Concentrated Flow, Segment #6
					Woodland Kv= 5.0 fps
0.1	40	0.0620	5.05		Shallow Concentrated Flow, Segment #7
					Paved Kv= 20.3 fps
4.8	370	0.0655	1.28		Shallow Concentrated Flow, Segment #8
					Woodland Kv= 5.0 fps
0.6	84	0.1165	2.39		Shallow Concentrated Flow, Segment #9
					Short Grass Pasture Kv= 7.0 fps
1.6	165	0.1150	1.70		Shallow Concentrated Flow, Segment #10
					Woodland Kv= 5.0 fps
8.4	622	0.0611	1.24		Shallow Concentrated Flow, Segment #11
					Woodland Kv= 5.0 fps
3.3	171	0.0293	0.86		Shallow Concentrated Flow, Segment #12
					Woodland Kv= 5.0 fps
71.7	1,862	0.0075	0.43		Shallow Concentrated Flow, Segment #13
					Woodland Kv= 5.0 fps
104.1	1,249	0.0016	0.20		Shallow Concentrated Flow, Segment #14
					Woodland Kv= 5.0 fps
16.4	519	0.0111	0.53		Shallow Concentrated Flow, Segment #15
					Woodland Kv= 5.0 fps
0.3	30	0.0083	1.85		Shallow Concentrated Flow, Segment #16
	400	0.0440	0.70		Paved Kv= 20.3 fps
4.1	182	0.0110	0.73		Shallow Concentrated Flow, Segment 317
4.4	0.4	0.0047	4.40		Short Grass Pasture Kv= 7.0 fps
1.4	91	0.0247	1.10		Shallow Concentrated Flow, Segment #18
40.4	004	0.0077	0.00		Short Grass Pasture Kv= 7.0 fps
12.1	604	0.0277	0.83		Shallow Concentrated Flow, Segment #19
40.0	000	0.0050	4.40		Woodland Kv= 5.0 fps
13.2	888	0.0056	1.12		Shallow Concentrated Flow, Segment #20
42.0	070	0.0064	4 47		Grassed Waterway Kv= 15.0 fps
13.9	978	0.0061	1.17		Shallow Concentrated Flow, Segment #21
0.6	740	0.0004	4 40		Grassed Waterway Kv= 15.0 fps
8.6	740	0.0091	1.43		Shallow Concentrated Flow, Segment #22
2.0	704	0.0720	4.00		Grassed Waterway Kv= 15.0 fps
2.9	704	0.0739	4.08		Shallow Concentrated Flow, Segment #23
	0.776	T			Grassed Waterway Kv= 15.0 fps
286.2	9,776	Total			

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 9

Subcatchment 1S: 1S



Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023

Page 10

Summary for Reach 100R: Final Reach #100

[40] Hint: Not Described (Outflow=Inflow)


 $Inflow\ Area = 1,263.040\ a\varsigma \quad 9.67\%\ Impervious,\ Inflow\ Depth > 1.31" \quad for\ 25Yr-24Hr\ event$


Inflow = 264.31 cfs @ 16.24 hrs, Volume= 137.379 af

Outflow = 264.31 cfs @ 16.24 hrs, Volume= 137.379 af, Atten= 0%, Lag= 0.0 min

Routing by Dyn-Stor-Ind method, Time Span= 0.00-24.00 hrs, dt= 0.05 hrs

Reach 100R: Final Reach #100





Type III 24-hr 2Yr-24Hr Rainfall=3.05"

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Page 2

Printed 9/19/2023

Time span=0.00-24.00 hrs, dt=0.05 hrs, 481 points
Runoff by SCS TR-20 method, UH=SCS, Weighted-CN
Reach routing by Dyn-Stor-Ind method - Pond routing by Dyn-Stor-Ind method

Subcatchment1S: 1S Runoff Area=55,018,008 sf 9.67% Impervious Runoff Depth>0.19"

Flow Length=9,776' Tc=286.2 min CN=56 Runoff=33.75 cfs 19.923 af

Reach100R: Final Reach#100Inflow=33.75 cfs 19.923 af
Outflow=33.75 cfs 19.923 af

Type III 24-hr 10Yr-24Hr Rainfall=4.60"

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 3

Time span=0.00-24.00 hrs, dt=0.05 hrs, 481 points
Runoff by SCS TR-20 method, UH=SCS, Weighted-CN
Reach routing by Dyn-Stor-Ind method - Pond routing by Dyn-Stor-Ind method

Subcatchment1S: 1S Runoff Area=55,018,008 sf 9.67% Impervious Runoff Depth>0.73"

Flow Length=9,776' Tc=286.2 min CN=56 Runoff=141.11 cfs 76.455 af

Reach100R: Final Reach#100 Inflow=141.11 cfs 76.455 af

Outflow=141.11 cfs 76.455 af

Type III 24-hr 25Yr-24Hr Rainfall=5.81"

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 4

Time span=0.00-24.00 hrs, dt=0.05 hrs, 481 points
Runoff by SCS TR-20 method, UH=SCS, Weighted-CN
Reach routing by Dyn-Stor-Ind method - Pond routing by Dyn-Stor-Ind method

Subcatchment1S: 1S Runoff Area=55,018,008 sf 9.67% Impervious Runoff Depth>1.31"

Flow Length=9,776' Tc=286.2 min CN=56 Runoff=264.31 cfs 137.379 af

Reach100R: Final Reach#100 Inflow=264.31 cfs 137.379 af

Outflow=264.31 cfs 137.379 af

Type III 24-hr 50Yr-24Hr Rainfall=6.94"

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 5

Time span=0.00-24.00 hrs, dt=0.05 hrs, 481 points
Runoff by SCS TR-20 method, UH=SCS, Weighted-CN
Reach routing by Dyn-Stor-Ind method - Pond routing by Dyn-Stor-Ind method

Subcatchment1S: 1S Runoff Area=55,018,008 sf 9.67% Impervious Runoff Depth>1.94"

Flow Length=9,776' Tc=286.2 min CN=56 Runoff=401.33 cfs 203.763 af

Reach100R: Final Reach#100 Inflow=401.33 cfs 203.763 af

Outflow=401.33 cfs 203.763 af

Type III 24-hr 100Yr-24Hr Rainfall=8.29"

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 6

Time span=0.00-24.00 hrs, dt=0.05 hrs, 481 points
Runoff by SCS TR-20 method, UH=SCS, Weighted-CN
Reach routing by Dyn-Stor-Ind method - Pond routing by Dyn-Stor-Ind method

Subcatchment1S: 1S Runoff Area=55,018,008 sf 9.67% Impervious Runoff Depth>2.78"

Flow Length=9,776' Tc=286.2 min CN=56 Runoff=583.92 cfs 292.122 af

Reach100R: Final Reach#100Inflow=583.92 cfs 292.122 af
Outflow=583.92 cfs 292.122 af

Appendix II - Proposed Conditions Analysis


25 Yr - 24 Hr. Full Summary 2 Yr - 24 Hr. Node Listing

10 Yr -24 Hr. Node Listing

25 Yr -24 Hr. Node Listing

50 Yr -24 Hr. Node Listing

100 Yr - 24 Hr. Node Listing



22-109 W-2 Proposed Analysis

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 2

Project Notes

CarlsonSurface||
RationalHydrographMethod|2|
ModifiedRational|5|1.00|1.00|
UnitHydrographMethod|1|
Rainfall events imported from "22-109 W-1 Existing Analysis.hcp"

Printed 9/19/2023 Page 3

Area Listing (all nodes)

Area	CN	Description
(acres)		(subcatchment-numbers)
46.159	39	>75% Grass cover, Good HSG A (1S)
22.856	61	>75% Grass cover, Good HSG B (1S)
44.540	74	>75% Grass cover, Good HSG C (1S)
8.799	80	>75% Grass cover, Good HSG D (1S)
5.027	76	Gravel roads HSG A (1S)
3.384	85	Gravel roads HSG B (1S)
1.789	89	Gravel roads HSG C (1S)
0.867	91	Gravel roads, HSG D (1S)
10.272	98	Paved parking HSG A (1S)
2.640	98	Paved parking HSG B (1S)
10.326	98	Paved parking HSG C (1S)
2.852	98	Paved parking HSG D (1S)
4.571	98	Roofs HSG A (1S)
1.003	98	Roofs HSG B (1S)
3.333	98	Roofs HSG C (1S)
0.758	98	Roofs HSG D (1S)
4.356	98	Water Surface HSG A (1S)
12.498	98	Water Surface HSG B (1S)
1.070	98	Water Surface HSG C (1S)
71.867	98	Water Surface HSG D (1S)
429.353	30	Woods, Good HSG A (1S)
256.991	55	Woods, Good HSG B (1S)
170.097	70	Woods, Good HSG C (1S)
147.633	77	Woods, Good HSG D (1S)
1,263.040	56	TOTAL AREA

22-109 W-2 Proposed AnalysisPrepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 4

Soil Listing (all nodes)

Area	Soil	Subcatchment
(acres)	Group	Numbers
499.737	HSG A	1S
299.372	HSG B	1S
231.154	HSG C	1S
232.776	HSG D	1S
0.000	Other	
1,263.040		TOTAL AREA

22-109 W-2 Proposed AnalysisPrepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 5

Ground Covers (all nodes)

HSG-/		HSG-C (acres)	HSG-D (acres)	Other (acres)	Total (acres)	Ground Cover	Subcatchment Numbers
46.15	9 22.856	44.540	8.799	0.000	122.354	>75% Grass cove	er, Good 1S
5.02	7 3.384	1.789	0.867	0.000	11.066	Gravel roads	1S
10.27	2 2.640	10.326	2.852	0.000	26.090	Paved parking	1S
4.57	1 1.003	3.333	0.758	0.000	9.664	Roofs	1S
4.35	6 12.498	1.070	71.867	0.000	89.791	Water Surface	1S
429.35	3 256.991	170.097	147.633	0.000	1,004.075	Woods, Good	1S
499.73	7 299.372	231.154	232.776	0.000	1,263.040	TOTAL AREA	

Type III 24-hr 25Yr-24Hr Rainfall=5.81"

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 6

Time span=0.00-24.00 hrs, dt=0.05 hrs, 481 points
Runoff by SCS TR-20 method, UH=SCS, Weighted-CN
Reach routing by Dyn-Stor-Ind method - Pond routing by Dyn-Stor-Ind method

Subcatchment1S: 1S Runoff Area=55,018,008 sf 9.94% Impervious Runoff Depth>1.31"

Flow Length=9,776' Tc=286.2 min CN=56 Runoff=264.31 cfs 137.379 af

Reach100R: Final Reach#100Inflow=264.31 cfs 137.379 af
Outflow=264.31 cfs 137.379 af

Total Runoff Area = 1,263.040 ac Runoff Volume = 137.379 af Average Runoff Depth = 1.31" 90.06% Pervious = 1,137.495 ac 9.94% Impervious = 125.545 ac

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023

Page 7

Summary for Subcatchment 1S: 1S

CarlsonPlanXYPos|1153158.4476|260306.7175|

Runoff = 264.31 cfs @ 16.24 hrs, Volume= 137.379 af, Depth> 1.31"

Runoff by SCS TR-20 method, UH=SCS, Weighted-CN, Time Span= 0.00-24.00 hrs, dt= 0.05 hrs Type III 24-hr 25Yr-24Hr Rainfall=5.81"

Area (sf)	CN	Description
199,092	98	Roofs HSG A
43,688	98	Roofs HSG B
145,176	98	Roofs HSG C
33,004	98	Roofs HSG D
2,010,665	39	>75% Grass cover, Good HSG A
995,608	61	>75% Grass cover, Good HSG B
1,940,170	74	>75% Grass cover, Good HSG C
383,294	80	>75% Grass cover, Good HSG D
447,434	98	Paved parking HSG A
115,020	98	Paved parking HSG B
449,781	98	Paved parking HSG C
124,234	98	Paved parking HSG D
18,702,622	30	Woods, Good HSG A
11,194,512	55	Woods, Good HSG B
7,409,442	70	Woods, Good HSG C
6,430,915	77	Woods, Good HSG D
189,750	98	Water Surface HSG A
544,416	98	Water Surface HSG B
46,599	98	Water Surface HSG C
3,130,536	98	Water Surface HSG D
218,969	76	Gravel roads HSG A
147,414	85	Gravel roads HSG B
77,919	89	Gravel roads HSG C
37,748	91	Gravel roads, HSG D
55,018,008	56	Weighted Average
49,549,278		90.06% Pervious Area
5,468,730		9.94% Impervious Area

22-109 W-2 Proposed AnalysisPrepared by Berry Surveying & Engineering

HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023


Page 8

Tc (min)	Length (feet)	Slope (ft/ft)	Velocity (ft/sec)	Capacity (cfs)	Description
14.2	100	0.0600	0.12		Sheet Flow, Segment #1
					Woods: Light underbrush n= 0.400 P2= 3.05"
8.0	59	0.0639	1.26		Shallow Concentrated Flow, Segment #2
					Woodland Kv= 5.0 fps
0.5	59	0.0679	1.82		Shallow Concentrated Flow, Segment #3
					Short Grass Pasture Kv= 7.0 fps
1.7	128	0.0644	1.27		Shallow Concentrated Flow, Segment #4
	40	0.0500	4.00		Woodland Kv= 5.0 fps
0.0	13	0.0583	4.90		Shallow Concentrated Flow, Segment #5
4 -	440	0.0057	4.00		Paved Kv= 20.3 fps
1.5	118	0.0657	1.28		Shallow Concentrated Flow, Segment #6
0.1	40	0.0620	5.05		Woodland Kv= 5.0 fps Shallow Concentrated Flow, Segment #7
0.1	40	0.0020	5.05		Paved Kv= 20.3 fps
4.8	370	0.0655	1.28		Shallow Concentrated Flow, Segment #8
4.0	370	0.0000	1.20		Woodland Kv= 5.0 fps
0.6	84	0.1165	2.39		Shallow Concentrated Flow, segment #9
0.0	01	0.1100	2.00		Short Grass Pasture Kv= 7.0 fps
1.6	165	0.1150	1.70		Shallow Concentrated Flow, Segment #10
					Woodland Kv= 5.0 fps
8.4	622	0.0611	1.24		Shallow Concentrated Flow, Segment #11
					Woodland Kv= 5.0 fps
3.3	171	0.0293	0.86		Shallow Concentrated Flow, Segment #12
					Woodland Kv= 5.0 fps
71.7	1,862	0.0075	0.43		Shallow Concentrated Flow, Segment #13
					Woodland Kv= 5.0 fps
104.1	1,249	0.0016	0.20		Shallow Concentrated Flow, Segment #14
					Woodland Kv= 5.0 fps
16.4	519	0.0111	0.53		Shallow Concentrated Flow, Segment #15
0.0	20	0.0000	4.05		Woodland Kv= 5.0 fps
0.3	30	0.0083	1.85		Shallow Concentrated Flow, Segment #16
4.1	100	0.0110	0.73		Paved Kv= 20.3 fps Shallow Concentrated Flow Segment #17
4.1	102	0.0110	0.73		Shallow Concentrated Flow, Segment #17 Short Grass Pasture Kv= 7.0 fps
1.4	91	0.0247	1.10		Shallow Concentrated Flow, Segment #18
1.7	31	0.0247	1.10		Short Grass Pasture Kv= 7.0 fps
12.1	604	0.0277	0.83		Shallow Concentrated Flow, Segment #19
	001	0.0277	0.00		Woodland Kv= 5.0 fps
13.2	888	0.0056	1.12		Shallow Concentrated Flow, Segment #20
					Grassed Waterway Kv= 15.0 fps
13.9	978	0.0061	1.17		Shallow Concentrated Flow, Segment #21
					Grassed Waterway Kv= 15.0 fps
8.6	740	0.0091	1.43		Shallow Concentrated Flow, Segment #22
					Grassed Waterway Kv= 15.0 fps
2.9	704	0.0739	4.08		Shallow Concentrated Flow, Segment #23
					Grassed Waterway Kv= 15.0 fps
286.2	9,776	Total			

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 9

Subcatchment 1S: 1S



Prepared by Berry Surveying & Engineering HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC Printed 9/19/2023

Page 10

Summary for Reach 100R: Final Reach #100


[40] Hint: Not Described (Outflow=Inflow)


Inflow Area = 1,263.040 ac, 9.94% Impervious, Inflow Depth > 1.31" for 25Yr-24Hr event Inflow = 264.31 cfs @ 16.24 hrs, Volume= 137.379 af

Outflow 264.31 cfs @ 16.24 hrs, Volume= 137.379 af, Atten= 0%, Lag= 0.0 min

Routing by Dyn-Stor-Ind method, Time Span= 0.00-24.00 hrs, dt= 0.05 hrs

Reach 100R: Final Reach #100





Type III 24-hr 2Yr-24Hr Rainfall=3.05"

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 2

Time span=0.00-24.00 hrs, dt=0.05 hrs, 481 points
Runoff by SCS TR-20 method, UH=SCS, Weighted-CN
Reach routing by Dyn-Stor-Ind method - Pond routing by Dyn-Stor-Ind method

Subcatchmentl S: 1SRunoff Area=55,018,008 sf 9.94% Impervious Runoff Depth>0.19"

Flow Length=9,776' Tc=286.2 min CN=56 Runoff=33.75 cfs 19.923 af

Reach100R: Final Reach#100Inflow=33.75 cfs 19.923 af
Outflow=33.75 cfs 19.923 af

Type III 24-hr 10Yr-24Hr Rainfall=4.60"

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 3

rage

Time span=0.00-24.00 hrs, dt=0.05 hrs, 481 points
Runoff by SCS TR-20 method, UH=SCS, Weighted-CN
Reach routing by Dyn-Stor-Ind method - Pond routing by Dyn-Stor-Ind method

Subcatchment1S: 1S Runoff Area=55,018,008 sf 9.94% Impervious Runoff Depth>0.73"

Flow Length=9,776' Tc=286.2 min CN=56 Runoff=141.11 cfs 76.455 af

Reach100R: Final Reach#100 Inflow=141.11 cfs 76.455 af

Outflow=141.11 cfs 76.455 af

Type III 24-hr 25Yr-24Hr Rainfall=5.81"

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Page 4

Printed 9/19/2023

Time span=0.00-24.00 hrs, dt=0.05 hrs, 481 points
Runoff by SCS TR-20 method, UH=SCS, Weighted-CN
Reach routing by Dyn-Stor-Ind method - Pond routing by Dyn-Stor-Ind method

Subcatchment1S: 1S Runoff Area=55,018,008 sf 9.94% Impervious Runoff Depth>1.31"

Flow Length=9,776' Tc=286.2 min CN=56 Runoff=264.31 cfs 137.379 af

Reach100R: Final Reach#100Inflow=264.31 cfs 137.379 af
Outflow=264.31 cfs 137.379 af

Type III 24-hr 50Yr-24Hr Rainfall=6.94"

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023 Page 5

Time span=0.00-24.00 hrs, dt=0.05 hrs, 481 points
Runoff by SCS TR-20 method, UH=SCS, Weighted-CN
Reach routing by Dyn-Stor-Ind method - Pond routing by Dyn-Stor-Ind method

Subcatchment1S: 1S Runoff Area=55,018,008 sf 9.94% Impervious Runoff Depth>1.94"

Flow Length=9,776' Tc=286.2 min CN=56 Runoff=401.33 cfs 203.763 af

Reach100R: Final Reach#100Inflow=401.33 cfs 203.763 af
Outflow=401.33 cfs 203.763 af

Type III 24-hr 100Yr-24Hr Rainfall=8.29"

Prepared by Berry Surveying & Engineering
HydroCAD® 10.00-25 s/n 10221 © 2019 HydroCAD Software Solutions LLC

Printed 9/19/2023

Page 6

Time span=0.00-24.00 hrs, dt=0.05 hrs, 481 points
Runoff by SCS TR-20 method, UH=SCS, Weighted-CN
Reach routing by Dyn-Stor-Ind method - Pond routing by Dyn-Stor-Ind method

Subcatchment1S: 1S Runoff Area=55,018,008 sf 9.94% Impervious Runoff Depth>2.78"

Flow Length=9,776' Tc=286.2 min CN=56 Runoff=583.92 cfs 292.122 af

Reach100R: Final Reach#100 Inflow=583.92 cfs 292.122 af

Outflow=583.92 cfs 292.122 af

Appendix III - Calculations, Charts, & Graphs

USGS StreamStats Extreme Precipitation Tables NCRS USDA Web-soil Map 2/7/23, 4:22 PM StreamStats


StreamStats Report

Region ID: NH

Workspace ID: NH20230207204759981000

Clicked Point (Latitude, Longitude): 43.21378, -71.03120

Time: 2023-02-07 15:48:25 -0500



Collapse All

> Basin Characteristics

Parameter Code	Parameter Description	Value	Unit
APRAVPRE	Mean April Precipitation	4.319	inches
CSL10_85	Change in elevation divided by length between points 10 and 85 percent of distance along main channel to basin divide - main channel method not known	63.7	feet per mi
DRNAREA	Area that drains to a point on a stream	1.93	square miles
WETLAND	Percentage of Wetlands	9.9214	percent

> Peak-Flow Statistics

Peak-Flow Statistics Parameters [Peak Flow Statewide SIR2008 5206]

Parameter Code	Parameter Name	Value	Units	Min Limit	Max Limit
DRNAREA	Drainage Area	1.93	square miles	0.7	1290
APRAVPRE	Mean April Precipitation	4.319	inches	2.79	6.23
WETLAND	Percent Wetlands	9.9214	percent	0	21.8
CSL10_85	Stream Slope 10 and 85 Method	63.7	feet per mi	5.43	543

Peak-Flow Statistics Flow Report [Peak Flow Statewide SIR2008 5206]

PII: Prediction Interval-Lower, Plu: Prediction Interval-Upper, ASEp: Average Standard Error of Prediction, SE: Standard Error (other -- see report)

Statistic	Value	Unit	PII	Plu	ASEp	Equiv. Yrs.
50-percent AEP flood	58.7	ft^3/s	36	95.8	30.1	3.2
20-percent AEP flood	103	ft^3/s	62.3	170	31.1	4.7

2/7/23, 4:22 PM StreamStats

Statistic	Value	Unit	PII	Plu	ASEp	Equiv. Yrs.	
10-percent AEP flood	140	ft^3/s	83	236	32.3	6.2	
4-percent AEP flood	193	ft^3/s	111	337	34.3	8	
2-percent AEP flood	237	ft^3/s	132	426	36.4	9	
1-percent AEP flood	291	ft^3/s	156	542	38.6	9.8	
0.2-percent AEP flood	425	ft^3/s	211	858	44.1	11	

Peak-Flow Statistics Citations

Olson, S.A.,2009, Estimation of flood discharges at selected recurrence intervals for streams in New Hampshire: U.S.Geological Survey Scientific Investigations Report 2008-5206, 57 p. (http://pubs.usgs.gov/sir/2008/5206/)

USGS Data Disclaimer: Unless otherwise stated, all data, metadata and related materials are considered to satisfy the quality standards relative to the purpose for which the data were collected. Although these data and associated metadata have been reviewed for accuracy and completeness and approved for release by the U.S. Geological Survey (USGS), no warranty expressed or implied is made regarding the display or utility of the data for other purposes, nor on all computer systems, nor shall the act of distribution constitute any such warranty.

USGS Software Disclaimer: This software has been approved for release by the U.S. Geological Survey (USGS). Although the software has been subjected to rigorous review, the USGS reserves the right to update the software as needed pursuant to further analysis and review. No warranty, expressed or implied, is made by the USGS or the U.S. Government as to the functionality of the software and related material nor shall the fact of release constitute any such warranty. Furthermore, the software is released on condition that neither the USGS nor the U.S. Government shall be held liable for any damages resulting from its authorized or unauthorized use.

USGS Product Names Disclaimer: Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Application Version: 4.12.0

StreamStats Services Version: 1.2.22

NSS Services Version: 2.2.1

Extreme Precipitation Tables

Northeast Regional Climate Center

Data represents point estimates calculated from partial duration series. All precipitation amounts are displayed in inches.

Smoothing Yes

State New Hampshire

Location

Longitude 71.036 degrees West **Latitude** 43.211 degrees North

Elevation 0 feet

Date/Time Tue, 04 Oct 2022 09:07:02 -0400

Extreme Precipitation Estimates

	5min	10min	15min	30min	60min	120min		1hr	2hr	3hr	6hr	12hr	24hr	48hr		1day	2day	4day	7day	10day	
1yr	0.26	0.40	0.49	0.65	0.81	1.02	1yr	0.70	0.98	1.19	1.53	1.96	2.54	2.79	1yr	2.25	2.69	3.10	3.82	4.39	1yr
2yr	0.32	0.49	0.61	0.80	1.01	1.28	2yr	0.87	1.16	1.48	1.88	2.39	3.05	3.40	2yr	2.70	3.27	3.77	4.49	5.12	2yr
5yr	0.37	0.57	0.72	0.96	1.23	1.57	5yr	1.06	1.44	1.84	2.35	3.01	3.85	4.35	5yr	3.41	4.18	4.80	5.66	6.40	5yr
10yr	0.40	0.64	0.80	1.09	1.42	1.85	10yr	1.23	1.69	2.17	2.79	3.58	4.60	5.23	10yr	4.07	5.03	5.76	6.74	7.58	10yr
25yr	0.47	0.74	0.95	1.30	1.73	2.28	25yr	1.50	2.09	2.69	3.49	4.51	5.81	6.70	25yr	5.14	6.44	7.34	8.50	9.50	25yr
50yr	0.52	0.84	1.07	1.50	2.02	2.68	50yr	1.74	2.46	3.19	4.15	5.38	6.94	8.07	50yr	6.14	7.76	8.82	10.14	11.26	50yr
100yr	0.59	0.95	1.22	1.73	2.35	3.15	100yr	2.03	2.89	3.76	4.92	6.40	8.29	9.74	100yr	7.34	9.36	10.60	12.09	13.36	100yr
200yr	0.65	1.07	1.38	1.99	2.74	3.71	200yr	2.37	3.41	4.45	5.86	7.64	9.91	11.75	200yr	8.77	11.30	12.74	14.43	15.86	200yr
500yr	0.77	1.27	1.66	2.41	3.37	4.60	500yr	2.91	4.23	5.54	7.34	9.63	12.55	15.06	500yr	11.11	14.48	16.26	18.25	19.91	500yr

Lower Confidence Limits

	5min	10min	15min	30min	60min	120min		1hr	2hr	3hr	6hr	12hr	24hr	48hr		1day	2day	4day	7day	10day	
1yr	0.23	0.36	0.44	0.59	0.73	0.90	1yr	0.63	0.88	0.93	1.25	1.52	1.96	2.47	1yr	1.74	2.38	2.88	3.30	3.88	1yr
2yr	0.31	0.48	0.59	0.80	0.99	1.17	2yr	0.86	1.15	1.35	1.81	2.33	2.95	3.29	2yr	2.62	3.16	3.65	4.38	5.00	2yr
5yr	0.35	0.54	0.67	0.92	1.16	1.40	5yr	1.01	1.37	1.60	2.13	2.76	3.53	3.96	5yr	3.13	3.81	4.41	5.31	5.93	5yr
10yr	0.38	0.59	0.73	1.02	1.32	1.60	10yr	1.14	1.56	1.81	2.42	3.11	4.02	4.55	10yr	3.56	4.38	5.08	6.13	6.72	10yr
25yr	0.44	0.67	0.83	1.19	1.57	1.91	25yr	1.35	1.86	2.12	2.82	3.62	4.76	5.45	25yr	4.21	5.24	6.14	7.44	8.20	25yr
50yr	0.49	0.74	0.93	1.33	1.79	2.19	50yr	1.55	2.14	2.38	3.18	4.05	5.39	6.23	50yr	4.77	5.99	7.08	8.60	9.48	50yr
100yr	0.55	0.83	1.04	1.50	2.05	2.51	100yr	1.77	2.45	2.68	3.58	4.52	6.09	7.11	100yr	5.39	6.84	8.17	9.94	10.83	100yr
200yr	0.61	0.92	1.16	1.69	2.35	2.88	200yr	2.03	2.81	3.01	4.02	5.05	6.87	8.75	200yr	6.08	8.41	9.45	11.48	12.40	200yr
500yr	0.72	1.07	1.37	1.99	2.83	3.47	500yr	2.44	3.40	3.53	4.70	5.87	8.02	10.60	500yr	7.09	10.19	11.45	13.92	14.77	500yr

Upper Confidence Limits

	5min	10min	15min	30min	60min	120min		1hr	2hr	3hr	6hr	12hr	24hr	48hr		1day	2day	4day	7day	10day	
1yr	0.28	0.43	0.53	0.71	0.87	1.07	1yr	0.75	1.05	1.23	1.71	2.16	2.76	3.05	1yr	2.44	2.94	3.37	4.11	4.76	1yr
2yr	0.33	0.50	0.62	0.84	1.03	1.24	2yr	0.89	1.21	1.45	1.93	2.49	3.19	3.54	2yr	2.82	3.40	3.92	4.62	5.27	2yr
5yr	0.39	0.60	0.75	1.03	1.31	1.57	5yr	1.13	1.53	1.83	2.46	3.14	4.19	4.73	5yr	3.70	4.55	5.20	6.01	6.86	5yr
10yr	0.46	0.70	0.87	1.22	1.57	1.90	10yr	1.36	1.86	2.20	2.98	3.78	5.18	5.91	10yr	4.59	5.68	6.46	7.34	8.38	10yr
25yr	0.56	0.85	1.06	1.51	1.98	2.44	25yr	1.71	2.39	2.83	3.87	4.84	6.89	7.95	25yr	6.10	7.64	8.60	9.64	10.65	25yr
50yr	0.65	0.98	1.22	1.76	2.37	2.94	50yr	2.05	2.88	3.43	4.69	5.85	8.55	9.97	50yr	7.57	9.59	10.70	11.81	12.98	50yr
100yr	0.76	1.14	1.43	2.07	2.83	3.55	100yr	2.44	3.47	4.16	5.72	7.09	10.63	12.51	100yr	9.40	12.03	13.30	14.49	15.82	100yr
200yr	0.88	1.32	1.68	2.43	3.38	4.29	200yr	2.92	4.19	5.05	6.97	8.58	13.25	14.90	200yr	11.73	14.33	16.53	17.77	19.32	200yr
500yr	1.08	1.60	2.06	3.00	4.26	5.50	500yr	3.68	5.37	6.50	9.08	11.07	17.77	19.94	500yr	15.73	19.18	22.04	23.31	25.19	500yr






NRCS

Natural Resources Conservation Service A product of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local participants

Custom Soil Resource Report for Strafford County, New Hampshire



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (https://offices.sc.egov.usda.gov/locator/app?agency=nrcs) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2 053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	
How Soil Surveys Are Made	
Soil Map	
Soil Map	
Legend	.11
Map Unit Legend	. 12
Map Unit Descriptions	. 14
Strafford County, New Hampshire	16
AcB—Acton fine sandy loam, 0 to 8 percent slopes	. 16
AdB—Acton very stony fine sandy loam, 0 to 8 percent slopes	.17
AdC—Acton very stony fine sandy loam, 8 to 15 percent slopes	.18
CfB—Charlton fine sandy loam, 3 to 8 percent slopes	.19
CfC—Charlton fine sandy loam, 8 to 15 percent slopes	. 20
CsB—Charlton fine sandy loam, 3 to 8 percent slopes, very stony	. 22
CsC—Charlton fine sandy loam, 8 to 15 percent slopes, very stony	
CsD—Charlton very stony fine sandy loam, 15 to 25 percent slopes	. 26
CvD—Charlton extremely stony fine sandy loam, 8 to 25 percent slopes	
Fa—Fresh water marsh	
GIB—Gloucester fine sandy loam, 3 to 8 percent slopes	.29
GIC—Gloucester fine sandy loam, 8 to 15 percent slopes	.30
GsB—Gloucester very stony fine sandy loam, 3 to 8 percent slopes	
GsC—Gloucester very stony fine sandy loam, 8 to 15 percent slopes	
HaB—Hinckley loamy sand, 3 to 8 percent slopes	. 33
HaC—Hinckley loamy sand, 8 to 15 percent slopes	. 35
HbE—Hinckley loamy sand, 15 to 60 percent slopes	. 37
HcC—Hollis-Charlton fine sandy loams, 8 to 15 percent slopes	.38
HdB—Hollis-Charlton very rocky fine sandy loams, 3 to 8 percent	
slopes	.40
HdC—Hollis-Charlton very rocky fine sandy loams, 8 to 15 percent	
slopes	.42
HfC—Hollis-Gloucester fine sandy loams, 8 to 15 percent slopes	.44
HgB—Hollis-Gloucester very rocky fine sandy loams, 3 to 8 percent	
slopes	.45
HgC—Hollis-Gloucester very rocky fine sandy loams, 8 to 15 percent	
slopes	.47
HgD—Hollis-Gloucester very rocky fine sandy loams, 15 to 25 percent	
slopes	.49
HID—Hollis-Gloucester extremely rocky fine sandy loams, 8 to 25	
percent slopes	
LeA—Leicester very stony fine sandy loam, 0 to 3 percent slopes	.52
LrA—Leicester-Ridgebury fine sandy loams, 0 to 3 percent slopes,	
very stony	.53
LrB—Leicester-Ridgebury fine sandy loams, 3 to 8 percent slopes,	
verv stonv.	.56

Custom Soil Resource Report

Mp—Freetown and Swansea mucky peats, 0 to 2 percent slopes	58
PbB—Paxton fine sandy loam, 3 to 8 percent slopes	60
PdC—Paxton fine sandy loam, 8 to 15 percent slopes, very stony	62
RgB—Ridgebury fine sandy loam, 3 to 8 percent slopes	63
RIB—Ridgebury fine sandy loam, 3 to 8 percent slopes, very stony	65
Sb—Saugatuck loamy sand	67
SuB—Sutton fine sandy loam, 0 to 8 percent slopes, very stony	68
W—Water	69
Wa—Whitman fine sandy loam, 0 to 3 percent slopes, very stony	70
WgB—Woodbridge fine sandy loam, 3 to 8 percent slopes	71
WsB—Woodbridge fine sandy loam, 0 to 8 percent slopes, very stony.	73
References	75

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

Custom Soil Resource Report

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and

Custom Soil Resource Report

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.



MAP LEGEND

Area of Interest (AOI)

Area of Interest (AOI)

Soils

Soil Map Unit Polygons

-

Soil Map Unit Lines

Soil Map Unit Points

Special Point Features

Blowout

Borrow Pit

Clay Spot

Closed Depression

Gravel Pit

Gravelly Spot

Landfill

A Lava Flow

Marsh or swamp

Mine or Quarry

Miscellaneous Water

Perennial Water

→ Saline Spot

Sandy Spot

Severely Eroded Spot

Sinkhole

Slide or Slip

Spoil Area

Stony Spot

m Ve

Very Stony Spot

87

Wet Spot Other

Δ

Special Line Features

Water Features

_

Streams and Canals

Transportation

+++ Rails

Interstate Highways

US Routes

Major Roads

Background

100

Aerial Photography

Local Roads

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:20.000.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service Web Soil Survey URL:

Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Strafford County, New Hampshire Survey Area Data: Version 23, Sep 9, 2022

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Jun 19, 2020—Sep 20, 2020

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
AcB	Acton fine sandy loam, 0 to 8 percent slopes	4.1	0.4%
AdB	Acton very stony fine sandy loam, 0 to 8 percent slopes	0.7	0.1%
AdC	Acton very stony fine sandy loam, 8 to 15 percent slopes	12.8	1.4%
CfB	Charlton fine sandy loam, 3 to 8 percent slopes	2.6	0.3%
CfC	Charlton fine sandy loam, 8 to 15 percent slopes	6.1	0.7%
CsB	Charlton fine sandy loam, 3 to 8 percent slopes, very stony	8.9	1.0%
CsC	Charlton fine sandy loam, 8 to 15 percent slopes, very stony	127.2	13.6%
CsD	Charlton very stony fine sandy loam, 15 to 25 percent slopes	25.7	2.7%
CvD	Charlton extremely stony fine sandy loam, 8 to 25 percent slopes	20.2	2.2%
Fa	Fresh water marsh	17.3	1.8%
GIB	Gloucester fine sandy loam, 3 to 8 percent slopes	11.6	1.2%
GIC	Gloucester fine sandy loam, 8 to 15 percent slopes	28.2	3.0%
GsB	Gloucester very stony fine sandy loam, 3 to 8 percent slopes	140.1	15.0%
GsC	Gloucester very stony fine sandy loam, 8 to 15 percent slopes	21.4	2.3%
НаВ	Hinckley loamy sand, 3 to 8 percent slopes	5.5	0.6%
HaC	Hinckley loamy sand, 8 to 15 percent slopes	0.2	0.0%
HbE	Hinckley loamy sand, 15 to 60 percent slopes	3.9	0.4%
HcC	Hollis-Charlton fine sandy loams, 8 to 15 percent slopes	2.7	0.3%
HdB	Hollis-Charlton very rocky fine sandy loams, 3 to 8 percent slopes	48.9	5.2%
HdC	Hollis-Charlton very rocky fine sandy loams, 8 to 15 percent slopes	14.1	1.5%

Custom Soil Resource Report

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
HfC	Hollis-Gloucester fine sandy loams, 8 to 15 percent slopes	6.2	0.7%
HgB	Hollis-Gloucester very rocky fine sandy loams, 3 to 8 percent slopes	23.4	2.5%
HgC	Hollis-Gloucester very rocky fine sandy loams, 8 to 15 percent slopes	131.6	14.1%
HgD	Hollis-Gloucester very rocky fine sandy loams, 15 to 25 percent slopes	35.1	3.8%
HID	Hollis-Gloucester extremely rocky fine sandy loams, 8 to 25 percent slopes	12.2	1.3%
LeA	Leicester very stony fine sandy loam, 0 to 3 percent slopes	5.5	0.6%
LrA	Leicester-Ridgebury fine sandy loams, 0 to 3 percent slopes, very stony	33.9	3.6%
LrB	Leicester-Ridgebury fine sandy loams, 3 to 8 percent slopes, very stony	5.6	0.6%
Мр	Freetown and Swansea mucky peats, 0 to 2 percent slopes	11.7	1.2%
PbB	Paxton fine sandy loam, 3 to 8 percent slopes	29.2	3.1%
PdC	Paxton fine sandy loam, 8 to 15 percent slopes, very stony	40.0	4.3%
RgB	Ridgebury fine sandy loam, 3 to 8 percent slopes	2.7	0.3%
RIB	Ridgebury fine sandy loam, 3 to 8 percent slopes, very stony	2.4	0.3%
Sb	Saugatuck loamy sand	13.2	1.4%
SuB	Sutton fine sandy loam, 0 to 8 percent slopes, very stony	2.3	0.2%
W	Water	19.4	2.1%
Wa	Whitman fine sandy loam, 0 to 3 percent slopes, very stony	38.8	4.2%
WgB	Woodbridge fine sandy loam, 3 to 8 percent slopes	9.9	1.1%
WsB	Woodbridge fine sandy loam, 0 to 8 percent slopes, very stony	9.5	1.0%
Totals for Area of Interest		934.8	100.0%

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas

Custom Soil Resource Report

shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An association is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Strafford County, New Hampshire

AcB—Acton fine sandy loam, 0 to 8 percent slopes

Map Unit Setting

National map unit symbol: 9d6j Elevation: 80 to 920 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Acton and similar soils: 85 percent Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Acton

Setting

Parent material: Till

Typical profile

H1 - 0 to 6 inches: fine sandy loam

H2 - 6 to 23 inches: very gravelly loamy sand

H3 - 23 to 42 inches: very cobbly loamy coarse sand

Properties and qualities

Slope: 0 to 8 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Moderately well drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00

in/hr)

Depth to water table: About 12 to 30 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 3.6 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2w

Hydrologic Soil Group: A/D

Ecological site: F144AY008CT - Moist Till Uplands

Hydric soil rating: No

Minor Components

Not named pan

Percent of map unit: 10 percent

Hydric soil rating: No

Leicester

Percent of map unit: 5 percent Landform: Depressions Hydric soil rating: Yes

AdB—Acton very stony fine sandy loam, 0 to 8 percent slopes

Map Unit Setting

National map unit symbol: 9d6k Elevation: 100 to 970 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Acton and similar soils: 85 percent Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Acton

Setting

Parent material: Till

Typical profile

H1 - 0 to 6 inches: very stony fine sandy loam H2 - 6 to 23 inches: very gravelly loamy sand

H3 - 23 to 42 inches: very cobbly loamy coarse sand

Properties and qualities

Slope: 0 to 8 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Moderately well drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00

in/hr)

Depth to water table: About 12 to 30 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 3.6 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: A/D

Ecological site: F144BY602ME - Sandy Toeslope

Hydric soil rating: No

Minor Components

Not named pan

Percent of map unit: 10 percent

Hydric soil rating: No

Leicester

Percent of map unit: 5 percent Landform: Depressions Hydric soil rating: Yes

AdC—Acton very stony fine sandy loam, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 9d6l Elevation: 130 to 970 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Acton and similar soils: 85 percent Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Acton

Settina

Parent material: Till

Typical profile

H1 - 0 to 6 inches: very stony fine sandy loam
H2 - 6 to 23 inches: very gravelly loamy sand

H3 - 23 to 42 inches: very cobbly loamy coarse sand

Properties and qualities

Slope: 8 to 15 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Moderately well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00

in/hr)

Depth to water table: About 12 to 30 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 3.6 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: A/D

Ecological site: F144BY602ME - Sandy Toeslope

Hydric soil rating: No

Minor Components

Not named pan

Percent of map unit: 10 percent

Hydric soil rating: No

Gloucester

Percent of map unit: 5 percent

Hydric soil rating: No

CfB—Charlton fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 2wh0n

Elevation: 0 to 1,440 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: All areas are prime farmland

Map Unit Composition

Charlton and similar soils: 85 percent Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Charlton

Setting

Landform: Ridges, hills, ground moraines

Landform position (two-dimensional): Summit, shoulder, backslope Landform position (three-dimensional): Nose slope, side slope, crest

Down-slope shape: Convex, linear Across-slope shape: Convex

Parent material: Coarse-loamy melt-out till derived from granite, gneiss, and/or

schist

Typical profile

Ap - 0 to 7 inches: fine sandy loam

Bw - 7 to 22 inches: gravelly fine sandy loam C - 22 to 65 inches: gravelly fine sandy loam

Properties and qualities

Slope: 3 to 8 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to high

(0.14 to 14.17 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Moderate (about 6.9 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2e

Hydrologic Soil Group: B

Ecological site: F144AY034CT - Well Drained Till Uplands

Hydric soil rating: No

Minor Components

Sutton

Percent of map unit: 8 percent Landform: Ground moraines, hills

Landform position (two-dimensional): Footslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Linear Hydric soil rating: No

Paxton

Percent of map unit: 5 percent

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Summit, shoulder, backslope

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Convex, linear Across-slope shape: Convex Hydric soil rating: No

Chatfield

Percent of map unit: 1 percent

Landform: Ridges, hills

Landform position (two-dimensional): Summit, shoulder, backslope Landform position (three-dimensional): Nose slope, side slope, crest

Down-slope shape: Convex

Across-slope shape: Linear, convex

Hydric soil rating: No

Leicester

Percent of map unit: 1 percent

Landform: Drainageways, depressions

Down-slope shape: Linear Across-slope shape: Concave

Hydric soil rating: Yes

CfC—Charlton fine sandy loam, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 2wh0q

Elevation: 0 to 1,440 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Charlton and similar soils: 85 percent *Minor components*: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Charlton

Setting

Landform: Ridges, hills, ground moraines
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope

Down-slope shape: Convex, linear Across-slope shape: Convex

Parent material: Coarse-loamy melt-out till derived from granite, gneiss, and/or

schist

Typical profile

Ap - 0 to 7 inches: fine sandy loam

Bw - 7 to 22 inches: gravelly fine sandy loam C - 22 to 65 inches: gravelly fine sandy loam

Properties and qualities

Slope: 8 to 15 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to high

(0.14 to 14.17 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Moderate (about 6.9 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: B

Ecological site: F144AY034CT - Well Drained Till Uplands

Hydric soil rating: No

Minor Components

Sutton, fine sandy loam

Percent of map unit: 5 percent

Landform: Ridges, hills, ground moraines
Landform position (two-dimensional): Footslope
Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Linear Hydric soil rating: No

Paxton

Percent of map unit: 5 percent

Landform: Hills, ground moraines, drumlins Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Convex, linear Across-slope shape: Convex Hydric soil rating: No

Chatfield

Percent of map unit: 3 percent

Landform: Ridges, hills

Landform position (two-dimensional): Summit, shoulder, backslope Landform position (three-dimensional): Nose slope, side slope, crest

Down-slope shape: Convex

Across-slope shape: Linear, convex

Hydric soil rating: No

Canton

Percent of map unit: 2 percent

Landform: Ridges, hills, ground moraines

Landform position (two-dimensional): Summit, shoulder, backslope Landform position (three-dimensional): Nose slope, side slope, crest

Down-slope shape: Convex, linear Across-slope shape: Convex

Hydric soil rating: No

CsB—Charlton fine sandy loam, 3 to 8 percent slopes, very stony

Map Unit Setting

National map unit symbol: 2wh0r

Elevation: 0 to 1,570 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Farmland of local importance

Map Unit Composition

Charlton, very stony, and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Charlton, Very Stony

Setting

Landform: Ridges, hills, ground moraines

Landform position (two-dimensional): Summit, shoulder, backslope Landform position (three-dimensional): Nose slope, side slope, crest

Down-slope shape: Convex, linear Across-slope shape: Convex

Parent material: Coarse-loamy melt-out till derived from granite, gneiss, and/or schist

Typical profile

Oe - 0 to 2 inches: moderately decomposed plant material

A - 2 to 4 inches: fine sandy loam

Bw - 4 to 27 inches: gravelly fine sandy loam C - 27 to 65 inches: gravelly fine sandy loam

Properties and qualities

Slope: 3 to 8 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to high

(0.14 to 14.17 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Moderate (about 8.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: B

Ecological site: F142XB009VT - Acidic Till Upland

Hydric soil rating: No

Minor Components

Sutton, very stony

Percent of map unit: 5 percent Landform: Hills, ground moraines

Landform position (two-dimensional): Footslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Linear Hydric soil rating: No

Paxton, very stony

Percent of map unit: 5 percent

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Summit, shoulder, backslope

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Convex, linear Across-slope shape: Convex

Hydric soil rating: No

Chatfield, very stony

Percent of map unit: 3 percent

Landform: Ridges, hills

Landform position (two-dimensional): Summit, shoulder, backslope Landform position (three-dimensional): Nose slope, side slope, crest

Down-slope shape: Convex

Across-slope shape: Linear, convex

Hydric soil rating: No

Leicester, very stony

Percent of map unit: 2 percent

Landform: Drainageways, depressions

Down-slope shape: Linear Across-slope shape: Concave

Hydric soil rating: Yes

CsC—Charlton fine sandy loam, 8 to 15 percent slopes, very stony

Map Unit Setting

National map unit symbol: 2wh0p

Elevation: 0 to 1,570 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Charlton, very stony, and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Charlton, Very Stony

Setting

Landform: Ridges, hills, ground moraines

Landform position (two-dimensional): Summit, shoulder, backslope

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Convex, linear Across-slope shape: Convex

Parent material: Coarse-loamy melt-out till derived from granite, gneiss, and/or

schist

Typical profile

Oe - 0 to 2 inches: moderately decomposed plant material

A - 2 to 4 inches: fine sandy loam

Bw - 4 to 27 inches: gravelly fine sandy loam C - 27 to 65 inches: gravelly fine sandy loam

Properties and qualities

Slope: 8 to 15 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to high

(0.14 to 14.17 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Moderate (about 8.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: B

Ecological site: F144AY034CT - Well Drained Till Uplands

Hydric soil rating: No

Minor Components

Sutton, very stony

Percent of map unit: 5 percent Landform: Hills, ground moraines

Landform position (two-dimensional): Footslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Linear Hydric soil rating: No

Paxton, very stony

Percent of map unit: 5 percent

Landform: Hills, drumlins, ground moraines

Landform position (two-dimensional): Summit, shoulder, backslope

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Convex, linear Across-slope shape: Convex Hydric soil rating: No

Chatfield, very stony

Percent of map unit: 3 percent

Landform: Ridges, hills

Landform position (two-dimensional): Summit, shoulder, backslope Landform position (three-dimensional): Nose slope, side slope, crest

Down-slope shape: Convex

Across-slope shape: Linear, convex

Hydric soil rating: No

Leicester, very stony

Percent of map unit: 2 percent

Landform: Hills, ground moraines, drainageways, depressions Landform position (two-dimensional): Footslope, toeslope

Landform position (three-dimensional): Base slope

Down-slope shape: Concave, linear Across-slope shape: Concave

Hydric soil rating: Yes

CsD—Charlton very stony fine sandy loam, 15 to 25 percent slopes

Map Unit Setting

National map unit symbol: 9d6w

Elevation: 0 to 1,000 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 120 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Charlton and similar soils: 85 percent *Minor components*: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Charlton

Setting

Parent material: Till

Typical profile

H1 - 0 to 13 inches: very stony fine sandy loam

H2 - 13 to 36 inches: fine sandy loam
H3 - 36 to 40 inches: gravelly loamy sand

Properties and qualities

Slope: 15 to 25 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00

in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 5.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: A

Ecological site: F144AY034CT - Well Drained Till Uplands

Hydric soil rating: No

Minor Components

Not named

Percent of map unit: 10 percent

Hollis

Percent of map unit: 5 percent Hydric soil rating: No

CvD—Charlton extremely stony fine sandy loam, 8 to 25 percent slopes

Map Unit Setting

National map unit symbol: 9d6x Elevation: 0 to 1,000 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 120 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Charlton and similar soils: 85 percent Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Charlton

Setting

Parent material: Till

Typical profile

H1 - 0 to 13 inches: extremely stony fine sandy loam

H2 - 13 to 36 inches: fine sandy loam H3 - 36 to 40 inches: gravelly loamy sand

Properties and qualities

Slope: 8 to 25 percent

Surface area covered with cobbles, stones or boulders: 9.0 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00

in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 5.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 7s

Hydrologic Soil Group: A

Ecological site: F144AY034CT - Well Drained Till Uplands

Minor Components

Not named

Percent of map unit: 5 percent Hydric soil rating: No

Hollis

Percent of map unit: 5 percent Hydric soil rating: No

Sutton

Percent of map unit: 3 percent Hydric soil rating: No

Rock outcrop

Percent of map unit: 2 percent Hydric soil rating: No

Fa—Fresh water marsh

Map Unit Setting

National map unit symbol: 9d72

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Fresh water marsh: 100 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Fresh Water Marsh

Setting

Landform: Marshes

Typical profile

H1 - 0 to 6 inches: mucky peat H2 - 6 to 16 inches: mucky peat H3 - 16 to 65 inches: variable

Properties and qualities

Slope: 0 to 1 percent

Drainage class: Very poorly drained

Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00

in/hr)

Depth to water table: About 0 inches Frequency of ponding: Frequent

Available water supply, 0 to 60 inches: Moderate (about 8.6 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 8w

Ecological site: F144BY210ME - Marsh Wetland Complex

Hydric soil rating: Yes

GIB—Gloucester fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 9d73 Elevation: 70 to 1,100 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Gloucester and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Gloucester

Setting

Parent material: Till

Typical profile

H1 - 0 to 14 inches: fine sandy loam

H2 - 14 to 28 inches: very gravelly loamy sand H3 - 28 to 40 inches: very gravelly coarse sand

Properties and qualities

Slope: 3 to 8 percent

Depth to restrictive feature: More than 80 inches Drainage class: Somewhat excessively drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): High to very high (6.00

to 20.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 3.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2s

Hydrologic Soil Group: A

Ecological site: F144AY032NH - Dry Till Uplands

Hydric soil rating: No

Minor Components

Not named pan

Percent of map unit: 5 percent

Hollis

Percent of map unit: 5 percent

Hydric soil rating: No

Acton

Percent of map unit: 5 percent

Hydric soil rating: No

GIC—Gloucester fine sandy loam, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 9d74

Elevation: 20 to 970 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Gloucester and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Gloucester

Settina

Parent material: Till

Typical profile

H1 - 0 to 14 inches: fine sandy loam

H2 - 14 to 28 inches: very gravelly loamy sand H3 - 28 to 40 inches: very gravelly coarse sand

Properties and qualities

Slope: 8 to 15 percent

Depth to restrictive feature: More than 80 inches Drainage class: Somewhat excessively drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): High to very high (6.00

to 20.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 3.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: A

Ecological site: F144AY032NH - Dry Till Uplands

Minor Components

Acton

Percent of map unit: 5 percent Hydric soil rating: No

Not named pan

Percent of map unit: 5 percent

Hydric soil rating: No

Hollis

Percent of map unit: 5 percent Hydric soil rating: No

GsB—Gloucester very stony fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 9d75 Elevation: 30 to 1,260 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Gloucester and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Gloucester

Setting

Parent material: Till

Typical profile

H1 - 0 to 14 inches: very stony fine sandy loam H2 - 14 to 28 inches: very gravelly loamy sand H3 - 28 to 40 inches: very gravelly coarse sand

Properties and qualities

Slope: 3 to 8 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches Drainage class: Somewhat excessively drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): High to very high (6.00

to 20.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 3.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: A

Ecological site: F144BY601ME - Dry Sand

Hydric soil rating: No

Minor Components

Acton

Percent of map unit: 5 percent

Hydric soil rating: No

Not named

Percent of map unit: 5 percent

Hydric soil rating: No

Hollis

Percent of map unit: 5 percent

Hydric soil rating: No

GsC—Gloucester very stony fine sandy loam, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 9d76

Elevation: 0 to 1.440 feet

Mean annual precipitation: 36 to 71 inches

Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Gloucester and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Gloucester

Setting

Parent material: Till

Typical profile

H1 - 0 to 14 inches: very stony fine sandy loam H2 - 14 to 28 inches: very gravelly loamy sand H3 - 28 to 40 inches: very gravelly coarse sand

Properties and qualities

Slope: 8 to 15 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches Drainage class: Somewhat excessively drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): High to very high (6.00

to 20.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 3.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: A

Ecological site: F144BY601ME - Dry Sand

Hydric soil rating: No

Minor Components

Not named

Percent of map unit: 5 percent

Hydric soil rating: No

Hollis

Percent of map unit: 5 percent

Hydric soil rating: No

Acton

Percent of map unit: 5 percent

Hydric soil rating: No

HaB—Hinckley loamy sand, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 2svm8

Elevation: 0 to 1,430 feet

Mean annual precipitation: 36 to 53 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 250 days

Farmland classification: Not prime farmland

Map Unit Composition

Hinckley and similar soils: 85 percent Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hinckley

Setting

Landform: Outwash deltas, outwash terraces, moraines, kames, outwash plains,

kame terraces, eskers

Landform position (two-dimensional): Summit, shoulder, backslope, footslope

Landform position (three-dimensional): Nose slope, side slope, base slope, crest,

riser, tread

Down-slope shape: Concave, convex, linear Across-slope shape: Convex, linear, concave

Parent material: Sandy and gravelly glaciofluvial deposits derived from gneiss

and/or granite and/or schist

Typical profile

Oe - 0 to 1 inches: moderately decomposed plant material

A - 1 to 8 inches: loamy sand

Bw1 - 8 to 11 inches: gravelly loamy sand Bw2 - 11 to 16 inches: gravelly loamy sand BC - 16 to 19 inches: very gravelly loamy sand

C - 19 to 65 inches: very gravelly sand

Properties and qualities

Slope: 3 to 8 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Excessively drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to very

high (1.42 to 99.90 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Very low (about 3.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3s

Hydrologic Soil Group: A

Ecological site: F144AY022MA - Dry Outwash

Hydric soil rating: No

Minor Components

Windsor

Percent of map unit: 8 percent

Landform: Outwash deltas, outwash terraces, moraines, kames, outwash plains, kame terraces. eskers

Landform position (two-dimensional): Summit, shoulder, backslope, footslope Landform position (three-dimensional): Nose slope, side slope, base slope, crest, riser, tread

Down-slope shape: Concave, convex, linear Across-slope shape: Convex, linear, concave

Hydric soil rating: No

Sudbury

Percent of map unit: 5 percent

Landform: Outwash deltas, outwash terraces, moraines, outwash plains, kame terraces

Landform position (two-dimensional): Backslope, footslope

Landform position (three-dimensional): Head slope, side slope, base slope, tread

Down-slope shape: Concave, linear Across-slope shape: Concave, linear

Agawam

Percent of map unit: 2 percent

Landform: Outwash deltas, outwash terraces, moraines, kames, outwash plains, kame terraces, eskers

Landform position (two-dimensional): Summit, shoulder, backslope, footslope Landform position (three-dimensional): Nose slope, side slope, base slope, crest, riser, tread

Down-slope shape: Concave, convex, linear Across-slope shape: Convex, linear, concave

Hydric soil rating: No

HaC—Hinckley loamy sand, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 2svm9

Elevation: 0 to 1,480 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Hinckley and similar soils: 85 percent Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hinckley

Setting

Landform: Outwash deltas, outwash terraces, moraines, kames, outwash plains,

kame terraces, eskers

Landform position (two-dimensional): Shoulder, backslope, footslope, toeslope Landform position (three-dimensional): Head slope, nose slope, side slope, crest,

Down-slope shape: Concave, convex, linear Across-slope shape: Convex, linear, concave

Parent material: Sandy and gravelly glaciofluvial deposits derived from gneiss and/or granite and/or schist

Typical profile

Oe - 0 to 1 inches: moderately decomposed plant material

A - 1 to 8 inches: loamy sand

Bw1 - 8 to 11 inches: gravelly loamy sand Bw2 - 11 to 16 inches: gravelly loamy sand BC - 16 to 19 inches: very gravelly loamy sand

C - 19 to 65 inches: very gravelly sand

Properties and qualities

Slope: 8 to 15 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Excessively drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to very

high (1.42 to 99.90 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Low (about 3.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: A

Ecological site: F144AY022MA - Dry Outwash

Hydric soil rating: No

Minor Components

Merrimac

Percent of map unit: 5 percent

Landform: Outwash terraces, moraines, kames, outwash plains, eskers

Landform position (two-dimensional): Shoulder, backslope, footslope, toeslope

Landform position (three-dimensional): Head slope, nose slope, side slope, crest,

riser

Down-slope shape: Convex Across-slope shape: Convex

Hydric soil rating: No

Windsor

Percent of map unit: 5 percent

Landform: Outwash deltas, outwash terraces, moraines, outwash plains, kame terraces, eskers, kames

Landform position (two-dimensional): Shoulder, backslope, footslope, toeslope Landform position (three-dimensional): Head slope, nose slope, side slope, crest,

Down-slope shape: Concave, convex, linear Across-slope shape: Convex, linear, concave

Hydric soil rating: No

Sudbury

Percent of map unit: 5 percent

Landform: Outwash deltas, outwash terraces, moraines, outwash plains, kame terraces

Landform position (two-dimensional): Backslope, footslope Landform position (three-dimensional): Base slope, tread

Down-slope shape: Concave, linear Across-slope shape: Concave, linear

HbE—Hinckley loamy sand, 15 to 60 percent slopes

Map Unit Setting

National map unit symbol: 2svmh

Elevation: 0 to 890 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 145 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Hinckley and similar soils: 85 percent Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hinckley

Setting

Landform: Outwash deltas, outwash terraces, moraines, kames, outwash plains,

kame terraces, eskers

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Nose slope, side slope, crest, head slope,

riser

Down-slope shape: Concave, convex, linear Across-slope shape: Convex, linear, concave

Parent material: Sandy and gravelly glaciofluvial deposits derived from gneiss

and/or granite and/or schist

Typical profile

Oe - 0 to 1 inches: moderately decomposed plant material

A - 1 to 8 inches: loamy sand

Bw1 - 8 to 11 inches: gravelly loamy sand Bw2 - 11 to 16 inches: gravelly loamy sand BC - 16 to 19 inches: very gravelly loamy sand

C - 19 to 65 inches: very gravelly sand

Properties and qualities

Slope: 15 to 60 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Excessively drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to very

high (1.42 to 99.90 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Low (about 3.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 7e

Hydrologic Soil Group: A

Ecological site: F144AY022MA - Dry Outwash

Hydric soil rating: No

Minor Components

Windsor

Percent of map unit: 10 percent

Landform: Outwash deltas, outwash terraces, kames, outwash plains, kame

terraces, eskers, moraines

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Head slope, nose slope, side slope, crest,

riser

Down-slope shape: Concave, convex, linear Across-slope shape: Convex, linear, concave

Hydric soil rating: No

Merrimac

Percent of map unit: 5 percent

Landform: Outwash terraces, moraines, kames, outwash plains, eskers

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Side slope, crest, head slope, nose slope,

riser

Down-slope shape: Concave, convex, linear Across-slope shape: Convex, linear, concave

Hydric soil rating: No

HcC—Hollis-Charlton fine sandy loams, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 9d7k

Elevation: 0 to 1,080 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 120 to 240 days

Farmland classification: Farmland of local importance

Map Unit Composition

Hollis and similar soils: 55 percent Charlton and similar soils: 35 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hollis

Setting

Parent material: Till

Typical profile

H1 - 0 to 14 inches: fine sandy loam H2 - 14 to 18 inches: bedrock

Properties and qualities

Slope: 8 to 15 percent

Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Drainage class: Well drained Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.60 to 6.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Very low (about 2.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: D

Ecological site: F144AY033MA - Shallow Dry Till Uplands

Hydric soil rating: No

Description of Charlton

Setting

Parent material: Till

Typical profile

H1 - 0 to 13 inches: fine sandy loam
H2 - 13 to 36 inches: fine sandy loam
H3 - 36 to 40 inches: gravelly loamy sand

Properties and qualities

Slope: 8 to 15 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00

in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 5.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: A

Ecological site: F144AY034CT - Well Drained Till Uplands

Hydric soil rating: No

Minor Components

Not named

Percent of map unit: 5 percent

Buxton

Percent of map unit: 5 percent Hydric soil rating: No

HdB—Hollis-Charlton very rocky fine sandy loams, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 9d7m

Elevation: 0 to 1,000 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 120 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Hollis and similar soils: 40 percent Charlton and similar soils: 30 percent Minor components: 30 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hollis

Setting

Parent material: Till

Typical profile

H1 - 0 to 14 inches: very stony fine sandy loam

H2 - 14 to 18 inches: bedrock

Properties and qualities

Slope: 3 to 8 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Drainage class: Well drained Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.60 to 6.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Very low (about 2.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: D

Ecological site: F144AY033MA - Shallow Dry Till Uplands

Description of Charlton

Setting

Parent material: Till

Typical profile

H1 - 0 to 13 inches: very stony fine sandy loam

H2 - 13 to 36 inches: fine sandy loam
H3 - 36 to 40 inches: gravelly loamy sand

Properties and qualities

Slope: 3 to 8 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00

in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 5.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: A

Ecological site: F144AY034CT - Well Drained Till Uplands

Hydric soil rating: No

Minor Components

Rock outcrop

Percent of map unit: 10 percent

Hydric soil rating: No

Leicester

Percent of map unit: 5 percent Landform: Depressions

Hydric soil rating: Yes

Sutton

Percent of map unit: 5 percent

Hydric soil rating: No

Buxton

Percent of map unit: 5 percent

Hydric soil rating: No

Not named

Percent of map unit: 5 percent

HdC—Hollis-Charlton very rocky fine sandy loams, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 9d7n Elevation: 0 to 1,200 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 120 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Hollis and similar soils: 40 percent Charlton and similar soils: 30 percent Minor components: 30 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hollis

Setting

Parent material: Till

Typical profile

H1 - 0 to 14 inches: very stony fine sandy loam

H2 - 14 to 18 inches: bedrock

Properties and qualities

Slope: 8 to 15 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Drainage class: Well drained Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.60 to 6.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Very low (about 2.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: D

Ecological site: F144AY033MA - Shallow Dry Till Uplands

Description of Charlton

Setting

Parent material: Till

Typical profile

H1 - 0 to 13 inches: very stony fine sandy loam

H2 - 13 to 36 inches: fine sandy loam
H3 - 36 to 40 inches: gravelly loamy sand

Properties and qualities

Slope: 8 to 15 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00

in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 5.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: A

Ecological site: F144AY034CT - Well Drained Till Uplands

Hydric soil rating: No

Minor Components

Rock outcrop

Percent of map unit: 10 percent

Hydric soil rating: No

Not named

Percent of map unit: 10 percent

Hydric soil rating: No

Woodbridge

Percent of map unit: 5 percent

Hydric soil rating: No

Sutton

Percent of map unit: 5 percent

HfC—Hollis-Gloucester fine sandy loams, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 9d7t Elevation: 30 to 1.120 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Hollis and similar soils: 55 percent Gloucester and similar soils: 35 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hollis

Setting

Parent material: Till

Typical profile

H1 - 0 to 14 inches: fine sandy loam H2 - 14 to 18 inches: bedrock

Properties and qualities

Slope: 8 to 15 percent

Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Drainage class: Well drained Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.60 to 6.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Very low (about 2.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: D

Ecological site: F144AY033MA - Shallow Dry Till Uplands

Hydric soil rating: No

Description of Gloucester

Setting

Parent material: Till

Typical profile

H1 - 0 to 14 inches: fine sandy loam

H2 - 14 to 28 inches: very gravelly loamy sand H3 - 28 to 40 inches: very gravelly coarse sand

Properties and qualities

Slope: 8 to 15 percent

Depth to restrictive feature: More than 80 inches Drainage class: Somewhat excessively drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): High to very high (6.00

to 20.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 3.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: A

Ecological site: F144AY032NH - Dry Till Uplands

Hydric soil rating: No

Minor Components

Not named

Percent of map unit: 7 percent

Hydric soil rating: No

Acton

Percent of map unit: 3 percent

Hydric soil rating: No

HgB—Hollis-Gloucester very rocky fine sandy loams, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 9d7v Elevation: 30 to 1,120 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Hollis and similar soils: 50 percent Gloucester and similar soils: 30 percent

Minor components: 20 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hollis

Setting

Parent material: Till

Typical profile

H1 - 0 to 14 inches: very stony fine sandy loam

H2 - 14 to 18 inches: bedrock

Properties and qualities

Slope: 3 to 8 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Drainage class: Well drained Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.60 to 6.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Very low (about 2.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: D

Ecological site: F144AY033MA - Shallow Dry Till Uplands

Hydric soil rating: No

Description of Gloucester

Settina

Parent material: Till

Typical profile

H1 - 0 to 14 inches: very stony fine sandy loam H2 - 14 to 28 inches: very gravelly loamy sand H3 - 28 to 40 inches: very gravelly coarse sand

Properties and qualities

Slope: 3 to 8 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches Drainage class: Somewhat excessively drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): High to very high (6.00

to 20.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 3.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: A

Ecological site: F144AY032NH - Dry Till Uplands

Hydric soil rating: No

Minor Components

Rock outcrop

Percent of map unit: 10 percent

Hydric soil rating: No

Not named

Percent of map unit: 5 percent

Hydric soil rating: No

Acton

Percent of map unit: 3 percent

Hydric soil rating: No

Leicester

Percent of map unit: 2 percent

Landform: Depressions Hydric soil rating: Yes

HgC—Hollis-Gloucester very rocky fine sandy loams, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 9d7w Elevation: 30 to 1,120 feet

Mean annual precipitation: 36 to 71 inches

Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Hollis and similar soils: 50 percent Gloucester and similar soils: 30 percent

Minor components: 20 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hollis

Setting

Parent material: Till

Typical profile

H1 - 0 to 14 inches: very stony fine sandy loam

H2 - 14 to 18 inches: bedrock

Properties and qualities

Slope: 8 to 15 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Drainage class: Well drained Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.60 to 6.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Very low (about 2.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: D

Ecological site: F144AY033MA - Shallow Dry Till Uplands

Hydric soil rating: No

Description of Gloucester

Setting

Parent material: Till

Typical profile

H1 - 0 to 14 inches: very stony fine sandy loam H2 - 14 to 28 inches: very gravelly loamy sand H3 - 28 to 40 inches: very gravelly coarse sand

Properties and qualities

Slope: 8 to 15 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches Drainage class: Somewhat excessively drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): High to very high (6.00

to 20.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 3.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: A

Ecological site: F144AY032NH - Dry Till Uplands

Hydric soil rating: No

Minor Components

Rock outcrop

Percent of map unit: 10 percent

Hydric soil rating: No

Not named

Percent of map unit: 7 percent

Acton

Percent of map unit: 3 percent Hydric soil rating: No

HgD—Hollis-Gloucester very rocky fine sandy loams, 15 to 25 percent slopes

Map Unit Setting

National map unit symbol: 9d7x Elevation: 0 to 1,230 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Hollis and similar soils: 50 percent Gloucester and similar soils: 30 percent

Minor components: 20 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hollis

Setting

Parent material: Till

Typical profile

H1 - 0 to 14 inches: very stony fine sandy loam

H2 - 14 to 18 inches: bedrock

Properties and qualities

Slope: 15 to 25 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Drainage class: Well drained Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.60 to 6.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Very low (about 2.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: D

Ecological site: F144AY033MA - Shallow Dry Till Uplands

Description of Gloucester

Setting

Parent material: Till

Typical profile

H1 - 0 to 14 inches: very stony fine sandy loam H2 - 14 to 28 inches: very gravelly loamy sand H3 - 28 to 40 inches: very gravelly coarse sand

Properties and qualities

Slope: 15 to 25 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches Drainage class: Somewhat excessively drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): High to very high (6.00

to 20.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 3.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: A

Ecological site: F144AY032NH - Dry Till Uplands

Hydric soil rating: No

Minor Components

Rock outcrop

Percent of map unit: 10 percent

Hydric soil rating: No

Not named

Percent of map unit: 10 percent

Hydric soil rating: No

HID—Hollis-Gloucester extremely rocky fine sandy loams, 8 to 25 percent slopes

Map Unit Setting

National map unit symbol: 9d7y Elevation: 100 to 1,610 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Hollis and similar soils: 40 percent Gloucester and similar soils: 25 percent

Minor components: 35 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hollis

Setting

Parent material: Till

Typical profile

H1 - 0 to 14 inches: extremely stony fine sandy loam

H2 - 14 to 18 inches: bedrock

Properties and qualities

Slope: 8 to 25 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Drainage class: Well drained Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.60 to 6.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Very low (about 2.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: D

Ecological site: F144BY702ME - Shallow and Moderately-deep Till

Hydric soil rating: No

Description of Gloucester

Setting

Parent material: Till

Typical profile

H1 - 0 to 14 inches: extremely stony fine sandy loam H2 - 14 to 28 inches: very gravelly loamy sand H3 - 28 to 40 inches: very gravelly coarse sand

Properties and qualities

Slope: 8 to 25 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches Drainage class: Somewhat excessively drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): High to very high (6.00

to 20.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 3.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: A

Ecological site: F144BY601ME - Dry Sand

Hydric soil rating: No

Minor Components

Rock outcrop

Percent of map unit: 20 percent

Hydric soil rating: No

Not named

Percent of map unit: 10 percent

Hydric soil rating: No

Acton

Percent of map unit: 3 percent

Hydric soil rating: No

Leicester

Percent of map unit: 2 percent Landform: Depressions Hydric soil rating: Yes

LeA—Leicester very stony fine sandy loam, 0 to 3 percent slopes

Map Unit Setting

National map unit symbol: 9d81

Elevation: 0 to 2.100 feet

Mean annual precipitation: 28 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 100 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Leicester and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Leicester

Setting

Landform: Depressions Parent material: Till

Typical profile

H1 - 0 to 5 inches: very stony fine sandy loam

H2 - 5 to 44 inches: gravelly fine sandy loam

Properties and qualities

Slope: 0 to 3 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Poorly drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.60 to 6.00 in/hr)

Depth to water table: About 0 to 12 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Low (about 5.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 7s

Hydrologic Soil Group: A/D

Ecological site: F144AY009CT - Wet Till Depressions

Hydric soil rating: Yes

Minor Components

Ridgebury

Percent of map unit: 5 percent Landform: Depressions Hydric soil rating: Yes

Not named wet

Percent of map unit: 5 percent Landform: Outwash terraces Hydric soil rating: Yes

Whitman

Percent of map unit: 5 percent Landform: Depressions Hydric soil rating: Yes

LrA—Leicester-Ridgebury fine sandy loams, 0 to 3 percent slopes, very stony

Map Unit Setting

National map unit symbol: 2xffr Elevation: 20 to 960 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 145 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Leicester, very stony, and similar soils: 60 percent Ridgebury, very stony, and similar soils: 30 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Leicester, Very Stony

Setting

Landform: Hills, ground moraines, drainageways, depressions Landform position (two-dimensional): Footslope, toeslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave, linear Across-slope shape: Concave

Parent material: Coarse-loamy melt-out till derived from gneiss, granite, and/or schist

Typical profile

Oe - 0 to 1 inches: moderately decomposed plant material

A - 1 to 7 inches: fine sandy loam

Bg - 7 to 18 inches: fine sandy loam

BC - 18 to 24 inches: fine sandy loam

C1 - 24 to 39 inches: gravelly fine sandy loam C2 - 39 to 65 inches: gravelly fine sandy loam

Properties and qualities

Slope: 0 to 3 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Poorly drained

Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to high

(0.14 to 14.17 in/hr)

Depth to water table: About 0 to 6 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water supply, 0 to 60 inches: High (about 9.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 5s

Hydrologic Soil Group: B/D

Ecological site: F144BY305ME - Wet Loamy Flat

Hydric soil rating: Yes

Description of Ridgebury, Very Stony

Setting

Landform: Hills, ground moraines, drumlins, drainageways, depressions

Landform position (two-dimensional): Footslope, toeslope Landform position (three-dimensional): Head slope, base slope

Down-slope shape: Concave Across-slope shape: Concave

Parent material: Coarse-loamy lodgment till derived from gneiss, granite, and/or

schist

Typical profile

Oe - 0 to 1 inches: moderately decomposed plant material

A - 1 to 6 inches: fine sandy loam Bw - 6 to 10 inches: sandy loam

Bg - 10 to 19 inches: gravelly sandy loam Cd - 19 to 66 inches: gravelly sandy loam

Properties and qualities

Slope: 0 to 3 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent Depth to restrictive feature: 15 to 35 inches to densic material

Drainage class: Poorly drained

Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately

low (0.00 to 0.14 in/hr)

Depth to water table: About 0 to 6 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Low (about 3.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 5s

Hydrologic Soil Group: D

Ecological site: F144BY305ME - Wet Loamy Flat

Hydric soil rating: Yes

Minor Components

Walpole

Percent of map unit: 5 percent

Landform: Outwash terraces, drainageways, depressions

Landform position (three-dimensional): Tread

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

Whitman, very stony

Percent of map unit: 3 percent

Landform: Hills, ground moraines, drumlins, drainageways, depressions

Landform position (two-dimensional): Toeslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

Swansea, mucky peat

Percent of map unit: 2 percent

Landform: Swamps, marshes, kettles, depressions, bogs

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

LrB—Leicester-Ridgebury fine sandy loams, 3 to 8 percent slopes, very stony

Map Unit Setting

National map unit symbol: 2xffs Elevation: 100 to 1,160 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 145 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Leicester, very stony, and similar soils: 60 percent Ridgebury, very stony, and similar soils: 30 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Leicester, Very Stony

Setting

Landform: Hills, ground moraines, drainageways, depressions Landform position (two-dimensional): Footslope, toeslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave, linear Across-slope shape: Concave

Parent material: Coarse-loamy melt-out till derived from gneiss, granite, and/or schist

Typical profile

Oe - 0 to 1 inches: moderately decomposed plant material

A - 1 to 7 inches: fine sandy loam
Bg - 7 to 18 inches: fine sandy loam
BC - 18 to 24 inches: fine sandy loam

C1 - 24 to 39 inches: gravelly fine sandy loam C2 - 39 to 65 inches: gravelly fine sandy loam

Properties and qualities

Slope: 3 to 8 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Poorly drained Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to high

(0.14 to 14.17 in/hr)

Depth to water table: About 0 to 6 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water supply, 0 to 60 inches: High (about 9.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: B/D

Ecological site: F144BY305ME - Wet Loamy Flat

Hydric soil rating: Yes

Description of Ridgebury, Very Stony

Setting

Landform: Hills, ground moraines, drumlins, drainageways, depressions

Landform position (two-dimensional): Footslope, toeslope Landform position (three-dimensional): Head slope, base slope

Down-slope shape: Concave Across-slope shape: Concave

Parent material: Coarse-loamy lodgment till derived from gneiss, granite, and/or

schist

Typical profile

Oe - 0 to 1 inches: moderately decomposed plant material

A - 1 to 6 inches: fine sandy loam Bw - 6 to 10 inches: sandy loam

Bg - 10 to 19 inches: gravelly sandy loam Cd - 19 to 66 inches: gravelly sandy loam

Properties and qualities

Slope: 3 to 8 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent Depth to restrictive feature: 15 to 35 inches to densic material

Drainage class: Poorly drained Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately

low (0.00 to 0.14 in/hr)

Depth to water table: About 0 to 6 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Low (about 3.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: D

Ecological site: F144BY305ME - Wet Loamy Flat

Hydric soil rating: Yes

Minor Components

Woodbridge, very stony

Percent of map unit: 5 percent

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Summit, backslope, footslope

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Convex Across-slope shape: Linear

Hydric soil rating: No

Walpole

Percent of map unit: 3 percent

Landform: Outwash terraces, drainageways, depressions

Landform position (three-dimensional): Tread

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

Whitman, very stony

Percent of map unit: 2 percent

Landform: Hills, ground moraines, drumlins, drainageways, depressions

Landform position (two-dimensional): Toeslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

Mp—Freetown and Swansea mucky peats, 0 to 2 percent slopes

Map Unit Setting

National map unit symbol: 2w68w

Elevation: 10 to 940 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 145 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Freetown and similar soils: 50 percent Swansea and similar soils: 30 percent

Minor components: 20 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Freetown

Setting

Landform: Marshes, kettles, depressions, swamps, bogs

Down-slope shape: Concave Across-slope shape: Concave

Parent material: Moderately decomposed organic material

Typical profile

Oe1 - 0 to 2 inches: mucky peat Oe2 - 2 to 79 inches: mucky peat

Properties and qualities

Slope: 0 to 2 percent

Surface area covered with cobbles, stones or boulders: 0.0 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Very poorly drained

Runoff class: Negligible

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to high

(0.14 to 14.17 in/hr)

Depth to water table: About 0 to 6 inches

Frequency of flooding: None Frequency of ponding: Frequent

Available water supply, 0 to 60 inches: Very high (about 20.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 8

Hydrologic Soil Group: B/D

Ecological site: F144AY043MA - Acidic Organic Wetlands

Hydric soil rating: Yes

Description of Swansea

Setting

Landform: Swamps, marshes, depressions, bogs, kettles

Down-slope shape: Concave Across-slope shape: Concave

Parent material: Moderately decomposed organic material over sandy and gravelly

glaciofluvial deposits

Typical profile

Oe1 - 0 to 12 inches: mucky peat Oe2 - 12 to 25 inches: mucky peat

Cg - 25 to 79 inches: sand

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Very poorly drained

Runoff class: Negligible

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to high

(0.14 to 14.17 in/hr)

Depth to water table: About 0 to 6 inches

Frequency of flooding: None Frequency of ponding: Frequent

Available water supply, 0 to 60 inches: High (about 11.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 8

Hydrologic Soil Group: B/D

Ecological site: F144AY043MA - Acidic Organic Wetlands

Hydric soil rating: Yes

Minor Components

Natchaug

Percent of map unit: 10 percent

Landform: Depressions, depressions, depressions

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

Scarboro

Percent of map unit: 4 percent

Landform: Outwash deltas, outwash terraces, drainageways, depressions

Landform position (three-dimensional): Tread

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

Whitman

Percent of map unit: 4 percent Landform: Hills, depressions

Landform position (two-dimensional): Toeslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

Maybid

Percent of map unit: 2 percent Landform: Marine terraces

Landform position (three-dimensional): Tread

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

PbB—Paxton fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 2t2qp

Elevation: 0 to 1,570 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: All areas are prime farmland

Map Unit Composition

Paxton and similar soils: 80 percent Minor components: 20 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Paxton

Setting

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Summit, shoulder, backslope Landform position (three-dimensional): Nose slope, side slope, crest

Down-slope shape: Convex, linear Across-slope shape: Convex

Parent material: Coarse-loamy lodgment till derived from gneiss, granite, and/or schist

Typical profile

Ap - 0 to 8 inches: fine sandy loam
Bw1 - 8 to 15 inches: fine sandy loam
Bw2 - 15 to 26 inches: fine sandy loam
Cd - 26 to 65 inches: gravelly fine sandy loam

Properties and qualities

Slope: 3 to 8 percent

Depth to restrictive feature: 18 to 39 inches to densic material

Drainage class: Well drained Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately

low (0.00 to 0.14 in/hr)

Depth to water table: About 18 to 37 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Low (about 3.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2s

Hydrologic Soil Group: C

Ecological site: F144AY007CT - Well Drained Dense Till Uplands

Hydric soil rating: No

Minor Components

Woodbridge

Percent of map unit: 9 percent

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Summit, backslope, footslope

Landform position (three-dimensional): Side slope

Down-slope shape: Concave Across-slope shape: Linear Hydric soil rating: No

Ridgebury

Percent of map unit: 6 percent

Landform: Hills, ground moraines, drainageways, depressions
Landform position (two-dimensional): Toeslope, backslope, footslope
Landform position (three-dimensional): Base slope, head slope, dip

Down-slope shape: Concave
Across-slope shape: Concave

Hydric soil rating: Yes

Charlton

Percent of map unit: 5 percent

Landform: Hills

Down-slope shape: Linear Across-slope shape: Linear Hydric soil rating: No

PdC—Paxton fine sandy loam, 8 to 15 percent slopes, very stony

Map Unit Setting

National map unit symbol: 2w677

Elevation: 0 to 1,330 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Paxton, very stony, and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Paxton, Very Stony

Setting

Landform: Hills, ground moraines, drumlins Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Convex, linear Across-slope shape: Linear, convex

Parent material: Coarse-loamy lodgment till derived from gneiss, granite, and/or

schist

Typical profile

Oe - 0 to 2 inches: moderately decomposed plant material

A - 2 to 10 inches: fine sandy loam
Bw1 - 10 to 17 inches: fine sandy loam
Bw2 - 17 to 28 inches: fine sandy loam
Cd - 28 to 67 inches: gravelly fine sandy loam

Properties and qualities

Slope: 8 to 15 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent Depth to restrictive feature: 20 to 43 inches to densic material

Drainage class: Well drained Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately

low (0.00 to 0.14 in/hr)

Depth to water table: About 18 to 37 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Low (about 4.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: C

Ecological site: F144AY007CT - Well Drained Dense Till Uplands

Hydric soil rating: No

Minor Components

Woodbridge, very stony

Percent of map unit: 8 percent

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Backslope, footslope

Landform position (three-dimensional): Side slope

Down-slope shape: Concave Across-slope shape: Linear Hydric soil rating: No

Charlton, very stony

Percent of map unit: 5 percent

Landform: Hills

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Convex Across-slope shape: Convex Hydric soil rating: No

Ridgebury, very stony

Percent of map unit: 2 percent

Landform: Hills, ground moraines, drumlins, drainageways, depressions

Landform position (two-dimensional): Footslope, toeslope Landform position (three-dimensional): Head slope, base slope

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

RgB—Ridgebury fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 2xffw

Elevation: 0 to 1,030 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Farmland of local importance

Map Unit Composition

Ridgebury and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Ridgebury

Setting

Landform: Hills, ground moraines, drumlins, drainageways, depressions

Landform position (two-dimensional): Footslope, toeslope Landform position (three-dimensional): Head slope, base slope

Down-slope shape: Concave Across-slope shape: Concave

Parent material: Coarse-loamy lodgment till derived from gneiss, granite, and/or

schist

Typical profile

Oe - 0 to 1 inches: moderately decomposed plant material

A - 1 to 6 inches: fine sandy loam Bw - 6 to 10 inches: sandy loam

Bg - 10 to 19 inches: gravelly sandy loam Cd - 19 to 66 inches: gravelly sandy loam

Properties and qualities

Slope: 3 to 8 percent

Depth to restrictive feature: 15 to 35 inches to densic material

Drainage class: Poorly drained

Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately

low (0.00 to 0.14 in/hr)

Depth to water table: About 0 to 6 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Low (about 3.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4w

Hydrologic Soil Group: D

Ecological site: F144AY009CT - Wet Till Depressions

Hydric soil rating: Yes

Minor Components

Woodbridge

Percent of map unit: 8 percent

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Summit, backslope, footslope

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Convex Across-slope shape: Linear Hydric soil rating: No

Scituate

Percent of map unit: 4 percent

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Summit, backslope, footslope

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Convex, linear Across-slope shape: Convex

Hydric soil rating: No

Whitman

Percent of map unit: 3 percent

Landform: Hills, ground moraines, drumlins, drainageways, depressions

Landform position (two-dimensional): Toeslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

RIB—Ridgebury fine sandy loam, 3 to 8 percent slopes, very stony

Map Unit Setting

National map unit symbol: 2xffx Elevation: 40 to 1,320 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Ridgebury, very stony, and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Ridgebury, Very Stony

Setting

Landform: Hills, ground moraines, drumlins, drainageways, depressions

Landform position (two-dimensional): Footslope, toeslope Landform position (three-dimensional): Head slope, base slope

Down-slope shape: Concave Across-slope shape: Concave

Parent material: Coarse-loamy lodgment till derived from gneiss, granite, and/or

schist

Typical profile

Oe - 0 to 1 inches: moderately decomposed plant material

A - 1 to 6 inches: fine sandy loam Bw - 6 to 10 inches: sandy loam

Bg - 10 to 19 inches: gravelly sandy loam Cd - 19 to 66 inches: gravelly sandy loam

Properties and qualities

Slope: 3 to 8 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent Depth to restrictive feature: 15 to 35 inches to densic material

Drainage class: Poorly drained

Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately

low (0.00 to 0.14 in/hr)

Depth to water table: About 0 to 6 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Low (about 3.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4w

Hydrologic Soil Group: D

Ecological site: F144AY009CT - Wet Till Depressions

Hydric soil rating: Yes

Minor Components

Woodbridge, very stony

Percent of map unit: 7 percent

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Summit, backslope, footslope

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Convex Across-slope shape: Linear Hydric soil rating: No

Whitman, very stony

Percent of map unit: 4 percent

Landform: Hills, ground moraines, drumlins, drainageways, depressions

Landform position (two-dimensional): Toeslope
Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

Walpole

Percent of map unit: 2 percent

Landform: Outwash terraces, drainageways, depressions

Landform position (three-dimensional): Tread

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

Scituate, very stony

Percent of map unit: 2 percent

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Summit, backslope, footslope

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Convex, linear Across-slope shape: Convex

Hydric soil rating: No

Sb—Saugatuck loamy sand

Map Unit Setting

National map unit symbol: 9d8r Elevation: 300 to 1,000 feet

Mean annual precipitation: 27 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 125 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Saugatuck and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Saugatuck

Setting

Landform: Outwash terraces Parent material: Outwash

Typical profile

H1 - 0 to 4 inches: loamy sand H2 - 4 to 7 inches: sand

H3 - 7 to 26 inches: loamy sand H4 - 26 to 42 inches: sand

Properties and qualities

Slope: 0 to 3 percent

Depth to restrictive feature: 10 to 16 inches to undefined

Drainage class: Poorly drained

Runoff class: High

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.60 to 2.00 in/hr)

Depth to water table: About 0 to 12 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Very low (about 1.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4w

Hydrologic Soil Group: B/D

Ecological site: F144AY028MA - Wet Outwash

Hydric soil rating: Yes

Minor Components

Not named wet

Percent of map unit: 15 percent

Landform: Outwash terraces Hydric soil rating: Yes

SuB—Sutton fine sandy loam, 0 to 8 percent slopes, very stony

Map Unit Setting

National map unit symbol: 2xfff Elevation: 0 to 1,410 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Sutton, very stony, and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Sutton, Very Stony

Setting

Landform: Hills, ground moraines

Landform position (two-dimensional): Footslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Linear

Parent material: Coarse-loamy melt-out till derived from gneiss, granite, and/or

schist

Typical profile

Oi - 0 to 2 inches: slightly decomposed plant material

A - 2 to 7 inches: fine sandy loam
Bw1 - 7 to 19 inches: fine sandy loam
Bw2 - 19 to 27 inches: sandy loam
C1 - 27 to 41 inches: gravelly sandy loam
C2 - 41 to 62 inches: gravelly sandy loam

Properties and qualities

Slope: 0 to 8 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Moderately well drained

Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to high

(0.14 to 14.17 in/hr)

Depth to water table: About 12 to 27 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm) Available water supply, 0 to 60 inches: Moderate (about 8.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: B/D

Ecological site: F144AY008CT - Moist Till Uplands

Hydric soil rating: No

Minor Components

Charlton, very stony

Percent of map unit: 7 percent

Landform: Ridges, hills, ground moraines

Landform position (two-dimensional): Summit, shoulder, backslope

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Convex, linear Across-slope shape: Convex Hydric soil rating: No

Canton, very stony

Percent of map unit: 4 percent Landform: Ridges, moraines, hills

Landform position (two-dimensional): Summit, shoulder, backslope

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Convex, linear Across-slope shape: Convex

Hydric soil rating: No

Leicester, very stony

Percent of map unit: 3 percent

Landform: Hills, ground moraines, drainageways, depressions

Landform position (two-dimensional): Footslope, toeslope

Landform position (three-dimensional): Base slope

Down-slope shape: Concave, linear Across-slope shape: Concave Hydric soil rating: Yes

.

Whitman, very stony

Percent of map unit: 1 percent

Landform: Hills, ground moraines, drumlins, drainageways, depressions

Landform position (two-dimensional): Toeslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

W-Water

Map Unit Composition

Water (less than 40 acres): 100 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Wa—Whitman fine sandy loam, 0 to 3 percent slopes, very stony

Map Unit Setting

National map unit symbol: 2zggn

Elevation: 130 to 970 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Whitman, very stony, and similar soils: 81 percent

Minor components: 19 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Whitman, Very Stony

Setting

Landform: Ground moraines, drumlins, drainageways, depressions, hills

Landform position (two-dimensional): Toeslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Concave

Parent material: Coarse-loamy lodgment till derived from granite and gneiss and/or

SUIISU

Typical profile

Oi - 0 to 1 inches: peat

A - 1 to 10 inches: fine sandy loam

Bg - 10 to 17 inches: gravelly fine sandy loam Cdg - 17 to 61 inches: fine sandy loam

Properties and qualities

Slope: 0 to 3 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent Depth to restrictive feature: 7 to 38 inches to densic material

Drainage class: Very poorly drained

Runoff class: Negligible

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately

low (0.00 to 0.14 in/hr)

Depth to water table: About 0 to 6 inches

Frequency of flooding: None Frequency of ponding: Frequent

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Low (about 3.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 5s

Hydrologic Soil Group: D

Ecological site: F144AY041MA - Very Wet Till Depressions

Hydric soil rating: Yes

Minor Components

Ridgebury, very stony

Percent of map unit: 10 percent

Landform: Hills, ground moraines, drumlins, drainageways, depressions

Landform position (two-dimensional): Footslope, toeslope Landform position (three-dimensional): Head slope, base slope

Down-slope shape: Concave Across-slope shape: Concave

Ecological site: F144AY009CT - Wet Till Depressions

Hydric soil rating: Yes

Scarboro

Percent of map unit: 5 percent

Landform: Outwash deltas, outwash terraces, depressions, drainageways

Landform position (three-dimensional): Tread

Down-slope shape: Concave Across-slope shape: Concave

Ecological site: F144AY031MA - Very Wet Outwash

Hydric soil rating: Yes

Swansea

Percent of map unit: 3 percent Landform: Swamps, marshes, bogs Down-slope shape: Concave Across-slope shape: Concave

Ecological site: F144AY043MA - Acidic Organic Wetlands

Hydric soil rating: Yes

Woodbridge, very stony

Percent of map unit: 1 percent

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Summit, backslope, footslope

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Concave Across-slope shape: Linear

Ecological site: F144AY037MA - Moist Dense Till Uplands

Hydric soil rating: No

WgB—Woodbridge fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 2t2ql Elevation: 0 to 1,470 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: All areas are prime farmland

Map Unit Composition

Woodbridge, fine sandy loam, and similar soils: 82 percent

Minor components: 18 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Woodbridge, Fine Sandy Loam

Setting

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Summit, backslope, footslope

Landform position (three-dimensional): Side slope

Down-slope shape: Concave Across-slope shape: Linear

Parent material: Coarse-loamy lodgment till derived from gneiss, granite, and/or

schist

Typical profile

Ap - 0 to 7 inches: fine sandy loam
Bw1 - 7 to 18 inches: fine sandy loam
Bw2 - 18 to 30 inches: fine sandy loam
Cd - 30 to 65 inches: gravelly fine sandy loam

Properties and qualities

Slope: 3 to 8 percent

Depth to restrictive feature: 20 to 39 inches to densic material

Drainage class: Moderately well drained

Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately

low (0.00 to 0.14 in/hr)

Depth to water table: About 18 to 30 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Low (about 3.6 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2w

Hydrologic Soil Group: C/D

Ecological site: F144AY037MA - Moist Dense Till Uplands

Hydric soil rating: No

Minor Components

Paxton

Percent of map unit: 10 percent

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Summit, shoulder, backslope Landform position (three-dimensional): Nose slope, side slope, crest

Down-slope shape: Convex, linear Across-slope shape: Convex Hydric soil rating: No

Ridgebury

Percent of map unit: 8 percent

Landform: Hills, ground moraines, drainageways, depressions

Landform position (two-dimensional): Toeslope, backslope, footslope Landform position (three-dimensional): Base slope, head slope, dip

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

WsB—Woodbridge fine sandy loam, 0 to 8 percent slopes, very stony

Map Unit Setting

National map unit symbol: 2t2qr Elevation: 0 to 1,440 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Woodbridge, very stony, and similar soils: 82 percent

Minor components: 18 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Woodbridge, Very Stony

Setting

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Summit, backslope, footslope

Landform position (three-dimensional): Side slope

Down-slope shape: Concave Across-slope shape: Linear

Parent material: Coarse-loamy lodgment till derived from gneiss, granite, and/or

schist

Typical profile

Oe - 0 to 2 inches: moderately decomposed plant material

A - 2 to 9 inches: fine sandy loam
Bw1 - 9 to 20 inches: fine sandy loam
Bw2 - 20 to 32 inches: fine sandy loam
Cd - 32 to 67 inches: gravelly fine sandy loam

Properties and qualities

Slope: 0 to 8 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent Depth to restrictive feature: 20 to 43 inches to densic material

Drainage class: Moderately well drained

Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately

low (0.00 to 0.14 in/hr)

Depth to water table: About 19 to 27 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm)

Available water supply, 0 to 60 inches: Low (about 4.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: C/D

Ecological site: F144AY037MA - Moist Dense Till Uplands

Hydric soil rating: No

Minor Components

Paxton, very stony

Percent of map unit: 10 percent

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Summit, shoulder, backslope

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Convex, linear Across-slope shape: Linear, convex

Hydric soil rating: No

Ridgebury, very stony

Percent of map unit: 8 percent

Landform: Hills, ground moraines, drumlins, drainageways, depressions

Landform position (two-dimensional): Toeslope

Landform position (three-dimensional): Head slope, base slope

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

References

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Federal Register. September 18, 2002. Hydric soils of the United States.

Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 054262

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 053577

Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 053580

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.


United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2 053374


United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/landuse/rangepasture/?cid=stelprdb1043084

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/scientists/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf





YOUNG ROAD, LLC 76 YOUNG ROAD 1.) OWNER: BARRINGTON, NH 03825

1A.) APPLICANT: PAUL THIBODEAU BARRINGTON, NH 03825

2.) TAX MAP 240, LOT 8

3.) LOT AREA: 2,855,458 Sq.Ft., 65.55 Ac.

4.) THE INTENT OF THIS PLAN IS TO DEMONSTRATE THE EXISTING DRAINAGE OVERVIEW OF THE LOCUS PARCEL.

5.) HYDROCAD USES A SERIES OF NODE SUFFIXES FOR NUMBERING PURPOSES (S=SUBCATCHMENT, P=POND DEVICE, R=REACH), TO SIMPLIFY ANNOTATION THESE SUFFIXES ARE LEFT OFF THE WATERSHED PLANS AND NODE TYPE IS DENOTED BY THE SYMBOL SHAPE ACCORDING TO THE DISPLAYED LEGEND WHICH COINCIDES WITH HYDROCAD GRAPHICS.

 $Ad\mathcal{C}$ - ACTON VERY STONY FINE SANDY LOAM, 8 TO 15%

CsC - CHARLTON FINE SANDY LOAM, 8 TO 15% SLOPES,

Fa - FRESH WATER MARSH

GIC - GLOUCESTER FINE SANDY LOAM, 8 TO 15% SLOPES

 ${\it GsB}$ - Gloucester very stony fine sandy loam, 8 to 15% slopes

HgC - HOLLIS-GLOUCESTER VERY ROCKY FINE SANDY LOAMS, 8 TO 15% SLOPES

 LrA - Leicester-Ridgebury fine sandy loams, 0 to 3% SLOPES, VERY STONY

SEE WEBSOIL USDA-NRCS

LEGEND:

— — 300— — EXISTING CONTOUR MAJOR STONE WALL ----- WETLAND LINE PROPERTY LINE _____


----- FLOW REACH

SOIL SERIES HfB NRCS SOIL LABEL LIMIT OF WATERSHED **■ ■ ■ ■ ■ ■** TIME OF CONCENTRATION PATH MATCH LINE

EXISTING CONTOUR MINOR


S.C.R.D. STRAFFORD COUNTY REGISTRY OF DEEDS
TYP. TYPICAL FND FOUND TBR TO BE REMOVED

SYMBOLS LEGEND:






SUBCATCHMENT AREA SYMBOL





POND DEVICE SYMBOL


> CONCENTRATION SEGMENT





SHEET 1 OF 10














YOUNG ROAD LLC 76 YOUNG ROAD 1.) OWNER: BARRINGTON, NH 03825

1A.) APPLICANT: PAUL THIBODEAU BARRINGTON, NH 03825

- 2.) TAX MAP 240, LOT 8
- 3.) LOT AREA: 2,855,458 Sq.Ft., 65.55 Ac.
- 4.) THE INTENT OF THIS PLAN IS TO DEMONSTRATE THE PROPOSED DRAINAGE OVERVIEW OF THE LOCUS PARCEL.
- 5.) HYDROCAD USES A SERIES OF NODE SUFFIXES FOR NUMBERING PURPOSES (S=SUBCATCHMENT, P=POND DEVICE, R=REACH), TO SIMPLIFY ANNOTATION THESE SUFFIXES ARE LEFT OFF THE WATERSHED PLANS AND NODE TYPE IS DENOTED BY THE SYMBOL SHAPE ACCORDING TO THE DISPLAYED LEGEND WHICH COINCIDES WITH HYDROCAD GRAPHICS.

 $Ad\mathcal{C}$ - ACTON VERY STONY FINE SANDY LOAM, 8 TO 15%

CsC - CHARLTON FINE SANDY LOAM, 8 TO 15% SLOPES,

Flpha - FRESH WATER MARSH

GIC - GLOUCESTER FINE SANDY LOAM, 8 TO 15% SLOPES

 ${\it GsB}$ - Gloucester very stony fine sandy loam, 8 to 15% slopes

HgC - HOLLIS-GLOUCESTER VERY ROCKY FINE SANDY LOAMS, 8 TO 15% SLOPES

 LrA - Leicester-Ridgebury fine sandy loams, 0 to 3% SLOPES, VERY STONY

SEE WEBSOIL USDA-NRCS


LEGEND:

_ _ _ _ _ _ 299 _ _ _ _ _ _ _ _ _ STONE WALL ______ 448A **—**----------

PROPERTY LINE SOIL LINE NRCS SOIL LINE NRCS SOIL LABEL LIMIT OF WATERSHED TIME OF CONCENTRATION PATH FLOW REACH MATCH LINE STRAFFORD COUNTY REGISTRY OF DEEDS


EXISTING CONTOUR MINOR EXISTING CONTOUR MAJOR


WETLAND LINE





SUBCATCHMENT AREA SYMBOL





CONCENTRATION


1 inch = 800 ft.


SHEET 6 OF 10

S.C.R.D. STRAFFORD COUNT TYP. TYPICAL FND FOUND TBR TO BE REMOVED

SYMBOLS LEGEND:







SEGMENT

